

A METHOD OF ACTIVATING T CELLS AND AGENTS USEFUL FOR SAME

## FIELD OF THE INVENTION

5 The present invention relates generally to a method of activating T cells and more particularly to a method of activating T cells using glycosylphosphatidylinositol (referred to herein as "GPI") molecules and derivatives or equivalents thereof. Even more particularly the method of the present invention contemplates a method of activating T cells, using GPI molecules, via a CD1-restricted pathway. The method of the present  
10 invention is useful, inter alia, in a range of therapeutic and/or prophylactic applications including, but not limited to applications which require skewing of the TH1/TH2 response or which require the induction of antibody production.

## BACKGROUND OF THE INVENTION

15

Bibliographic details of the publications numerically referred to in this specification are collected at the end of the description.

Antibody responses to protein are understood to be MHC restricted. That is, the  
20 production of antibodies directed to a given T dependent antigen requires the production of cytokines by stimulated TH2 cells. Said TH2 cells are stimulated following their binding to a MHC II/peptide complex comprising a peptide derived from the processing of said antigen. Since MHC molecules are polymorphic, there exist genetically determined high and low responders to peptide vaccines.

25

GPI anchor surface proteins occur frequently among medically important parasitic and fungal taxa such as *Plasmodium*, *Trypanosoma*, *Leishmania*, *Toxoplasma* and *Candida*.

GPIs are ubiquitous among eukaryotes, described from *T. brucei*, *T. cruzi*, *Plasmodium*,  
30 *Leishmania*, and *Toxoplasma*, as well as yeast, insect, fish and numerous mammalian sources (for recent reviews see (1, 2)). GPIs consist of a conserved core glycan (Man $\alpha$ 1-

09830451-073101

- 2 -

2Man $\alpha$ 1-6Man $\alpha$ 1-4GlcNH<sub>2</sub> linked to the 6-position of the *myo*-inositol ring of PI. GPIs are built up on the cytoplasmic face of the endoplasmic reticulum by the sequential addition of sugar residues to PI by the action of glycosyltransferases. The maturing GPI is then translocated across the membrane to the luminal side of the ER, whence it may be exported  
5 to the cell surface, free or in covalent association with proteins. The terasaccharide core glycan may be further substituted with sugars, phosphates and ethanolamine groups in a species and tissue-specific manner. GPI fatty acid moieties can be either diacylglycerols, alkylacylglycerols, monoalkylglycerols or ceramides, with additional palmitoylations or myristoylations to the inositol ring. The overall picture is of a closely related family of  
10 glycolipids sharing certain core features but with a high level of variation in fatty acid composition and side-chain modifications to the conserved core glycan.

In work leading up to the present invention, the inventors have shown that the antibody response to several parasitic proteins is regulated predominantly through CD1-restricted  
15 recognition of the covalently associated GPI moiety by IL-4 producing CD4<sup>+</sup> T cells with limited TCR repertoire diversity. In contrast, there is little evidence for MHC II restricted T cell responses to these antigens. GPI moieties, therefore, act as universal T cell sites through presentation by the non-polymorphic CD1 restriction element.

## 20 SUMMARY OF THE INVENTION

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated integer or group of integers but not the  
25 exclusion of any other integer or group of integers.

One aspect of the present invention contemplates a method of activating T cells, said method comprising administering a T cell activating effective amount of a molecule or a complex comprising said molecule which molecule or molecule complex is capable of  
30 interacting with CD1 on an immune cell to form an association with CD1 which association activates helper T cells.

09830451.073101

- 3 -

Another aspect of the present invention contemplates a method of activating T cells said method comprising administering a T cell activating effective amount of GPI or derivative or equivalent thereof or a complex comprising GPI or derivative or equivalent thereof which GPI or GPI-complex is capable of interacting with CD1 on an immune cell to form  
 5 an association with CD1 which association activates helper T cells.

Yet another aspect of the present invention contemplates a method of activating helper T cells said method comprising administering a T cell activating effective amount of *Plasmodium* GPI or derivative or equivalent thereof or a complex comprising said  
 10 *Plasmodium* GPI or derivative or equivalent thereof which *Plasmodium* GPI or *Plasmodium* GPI complex is capable of interacting with CD1 on an immune cell to form an association with CD1 which association activates helper T cells.

In still another aspect, said GPI comprises a structure selected from:

15

EtN-P-[M $\alpha$ 2]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y

EtN-P-[M $\alpha$ 2][G]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y

EtN-P-[M $\alpha$ 2][X]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y

EtN-P-[M $\alpha$ 2][EtN-P]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y

20

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 25 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In still yet another aspect, said GPI comprises a structure selected from:

30

EtN-P-M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y

09830451.073101

- 4 -

M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-YEtN-P-M $\alpha$ 2M $\alpha$ 6M-Y

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is  
 5 mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine  
 including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 linkages which may be substituted with any other positional linkages as required, and Y  
 10 is any lipid or phospholipid.

In yet still another aspect, said GPI comprises a structure selected from:

EtN-P-[M $\alpha$ 2][G]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y15 EtN-P-[M $\alpha$ 2][X]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-YEtN-P-[M $\alpha$ 2][EtN-P]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is  
 mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine  
 20 including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 linkages which may be substituted with any other positional linkages as required, and Y  
 is any lipid or phospholipid.

25

In a further aspect, said GPI comprises a structure selected from:

M $\alpha$ 2[M $\alpha$ 2][G]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-YM $\alpha$ 2[M $\alpha$ 2][X]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y30 M $\alpha$ 2[M $\alpha$ 2][EtN-P]M $\alpha$ 6M $\alpha$ 4G-Y

09830451-073101

- 5 -

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.


In another further aspect, said GPI comprises a structure selected from:

10



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In still another further aspect, said GPI comprises a structure selected from:



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may

09830451.073101

- 6 -

be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

5 In still yet another further aspect, said GPI comprises a structure selected from:

EtN-P-[M $\alpha$ 2][G]M $\alpha$ 2M $\alpha$ 6M-Y

EtN-P-[M $\alpha$ 2][X]M $\alpha$ 2M $\alpha$ 6M-Y

EtN-P-[M $\alpha$ 2][EtN-P]M $\alpha$ 2M $\alpha$ 6M-Y

10

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may

15 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In yet still another further aspect, said GPI comprises a structure selected from:

20

M $\alpha$ 2[M $\alpha$ 2][G]M $\alpha$ 2M $\alpha$ 6M-Y

M $\alpha$ 2[M $\alpha$ 2][X]M $\alpha$ 2M $\alpha$ 6M-Y

M $\alpha$ 2[M $\alpha$ 2][EtN-P]M $\alpha$ 6M-Y

25 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
30 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

09330451.073101

In still yet another further aspect, said GPI comprises a structure selected from:




5

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
10 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In another aspect, said GPI comprises a structure selected from:

15



20 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
25 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

Another aspect of the present invention contemplates a method of activating helper T cells said method comprising administering a T cell activating effective amount of GPI or  
30 derivative or equivalent thereof or a complex comprising said GPI or derivative or equivalent thereof which GPI or GPI complex is capable of interacting with CD1 on an

09830451-073101

immune cell to form an association with CD1 which association activates CD4<sup>+</sup> NK1.1<sup>+</sup> T cells.

Yet another aspect of the present invention contemplates a method of activating CD4<sup>+</sup>,  
5 NK1.1<sup>+</sup> T cells said method comprising administering a T cell activating effective amount  
of a *Plasmodium* GPI or derivative or equivalent thereof or a complex comprising said  
*Plasmodium* GPI or derivative or equivalent thereof which *Plasmodium* GPI or  
*Plasmodium* GPI complex is capable of interacting with CD1 on an immune cell to form  
an association with CD1 which association activates CD4<sup>+</sup> NK1.1<sup>+</sup> T cells.

10

Still another aspect of the present invention contemplates a method of activating T helper  
cells said method comprising administering a T cell activating effective amount of a GPI  
or derivative or equivalent thereof or a complex comprising said GPI or derivative or  
equivalent thereof which GPI or GPI complex is capable of interacting with CD1 on an  
15 immune cell to form an association with CD1 which association activates T helper cells  
wherein said activated T cells provide B cell help.

Still yet another aspect of the present invention contemplates a method of activating CD4<sup>+</sup>,  
NK1.1<sup>+</sup> T cells said method comprising administering a T cell activating effective amount  
20 of a *Plasmodium* GPI or derivative or equivalent thereof or a complex comprising said  
*Plasmodium* GPI or derivative or equivalent thereof which *Plasmodium* GPI or  
*Plasmodium* GPI complex is capable of interacting with CD1 on an immune cell to form  
an association with CD1 which association activates CD4<sup>+</sup>, NK1.1<sup>+</sup> T cells wherein said  
activated T cells provide B cell help.

25

Yet still another aspect of the present invention contemplates a method of activating T  
helper cells said method comprising administering a T cell activating effective amount of  
a GPI or derivative or equivalent thereof or a complex comprising said GPI or derivative  
or equivalent thereof which GPI or GPI complex is capable of interacting with CD1 on an  
30 immune cell to form an association with CD1 which association activates T helper cells  
wherein said activated T cells induce or otherwise upregulate a TH1-type response.

09330451-073101



A further aspect of the present invention contemplates a method of activating T helper cells said method comprising administering a T cell activating effective amount of a GPI or derivative or equivalent thereof or a complex comprising said GPI or derivative or  
5 equivalent thereof which GPI or GPI complex is capable of interacting with CD1 on an immune cell to form an association with CD1 which association activates T helper cells wherein said activated T cells induce or otherwise upregulate a TH2-type response.

Another further aspect of the present invention provides a method of inducing, in a  
10 mammal, an immune response directed to a GPI said method comprising administering to said mammal a T cell activating effective amount of GPI or derivative or equivalent thereof which GPI is capable of interacting with CD1 on an immune cell to form an association with CD1 which association activates helper T cells.

15 Still another further aspect of the present invention provides a method of inducing, in a mammal, an immune response directed to an antigen, said method comprising administering to said mammal a T cell activating effective amount of GPI or derivative or equivalent thereof complexed to said antigen, which GPI-antigen complex is capable of interacting with CD1 on an immune cell to form an association with CD1, which  
20 association activates helper T cells.

Still yet another further aspect of the present invention contemplates a method of treating a mammal said method comprising administering to said mammal an effective amount of GPI or derivative or equivalent thereof or a complex comprising said GPI or derivative or  
25 equivalent thereof which GPI or GPI complex is capable of interacting with CD1 on an immune cell to form an association with CD1 which association activates helper T cells.

Yet still another further aspect of the present invention contemplates a method of treating a mammal said method comprising administering a mammalian helper T cell activation  
30 effective amount of a GPI or derivative or equivalent thereof or a complex comprising said GPI or derivative or equivalent thereof which GPI or GPI complex is capable of interacting

- 10 -

with CD1 on an immune cell to form an association with CD1 which association activates helper T cells.

Another aspect of the present invention contemplates the use of GPI or derivative or  
5 equivalent thereof or a complex comprising said GPI or derivative or equivalent thereof  
which GPI or GPI complex is capable of interacting with CD1 on an immune cell, in the  
manufacture of a medicament for the activation of helper T cells in a mammal.

Yet another aspect of the present invention provides a method for the treatment and/or  
10 prophylaxis of a mammalian disease condition characterised by a micro-organism infection,  
said method comprising administering to said mammal an effective amount of GPI or  
derivative or equivalent thereof or a complex comprising said GPI or derivative or  
equivalent thereof which GPI or GPI complex is capable of interacting with CD1 on an  
immune cell to form an association with CD1 which association activates helper T cells.

15

Still another aspect of the present invention provides a method for the treatment and/or  
prophylaxis of a mammalian disease condition characterised by the insufficiency or absence  
of an appropriate TH1 response said method comprising administering to said mammal an  
effective amount of GPI or derivative or equivalent thereof or a complex comprising said  
20 GPI or derivative or equivalent thereof which GPI or GPI complex is capable of interacting  
with CD1 on an immune cell to form an association with CD1 which association induces  
or otherwise upregulates a TH1 response.

Still yet another aspect of the present invention provides a method for the treatment and/or  
25 prophylaxis of a mammalian disease condition characterised by the insufficiency or absence  
of an appropriate TH2 response said method comprising administering to said mammal an  
effective amount of GPI or derivative or equivalent thereof or a complex comprising said  
GPI or derivative or equivalent thereof which GPI or GPI complex is capable of interacting  
with CD1 on an immune cell to form an association with CD1 which association induces  
30 or otherwise upregulates a TH2 response.

- 11 -

Yet still another aspect of the present invention is directed to a composition which activates T cells, said composition comprising a GPI or derivative or equivalent thereof or a complex comprising GPI or derivative or equivalent thereof which GPI or GPI-complex is capable of interacting with CD1 on an immune cell to form an association with CD1  
5 which association activates helper T cells.

A further aspect of the present invention relates to an immunogenic composition comprising as the active component GPI or derivative or equivalent thereof or GPI complex, as broadly described above.

10

Another further aspect of the present invention is directed to a pharmaceutical composition capable of activating T cells, said composition comprising a GPI or derivative or equivalent thereof or a complex comprising GPI or derivative or equivalent thereof which GPI or GPI-complex is capable of interacting with CD1 on an immune cell to form an  
15 association with CD1, which association activates helper T cells, together with one or more pharmaceutically acceptable carriers and/or diluents.

20

09830451-073101  
T01E20-T540E860

### BRIEF DESCRIPTION OF THE DRAWINGS

**Figure 1** is a graphical representation of the survival from PbA infection in Balb/c mice wild type (○), Balb/c mice CD1d<sup>-/-</sup> (●), and C57B/6 (▲) mice. The results are from a single experiment representative of 2 independent experiments.

**Figure 2** is a graphical representation of the percentage of CD4<sup>+</sup> cells producing IL-4 or IFN- $\gamma$  during the infection with PbA. Balb/c wild type (□) and Balb/c CD1<sup>-/-</sup> (■) mice were infected with 1 x 10<sup>6</sup> PbA infected red blood cells. At day 7 post-infection the animals were sacrificed and the spleen cells stained for CD4<sup>+</sup> and intracellular cytokine content. The percentage of CD4<sup>+</sup> cells producing IL-4 or IFN- $\gamma$  is indicated.

**Figure 3.** Diagrammatical representation of GPI structures used in this study. Purification and compositional analyses are as described. (A) COOH-terminal GPIs from *T. brucei* and *P. falciparum*. Boxed areas represent modifications found in PfGPI. The cleavage site of mfVSG by phosphatidylinositol-specific phospholipase C (PI-PLC) is indicated. (B) Free iM2 iM4 and EP-iM4 GPIs of *L. mexicana*. Nomenclature is as described, where all isomers contain one mannose in  $\alpha$ -1-3 linkage, EP indicates ethanolamine phosphate, and M2, M4 indicate number of mannose residues, as shown. (C) Chemically synthesized rat brain Thy-1 GPI.

**Figure 4.** Response of peripheral NKT cells to purified GPIs *in vitro*. (A) As determined by forward (FSC) and side (SSC) light scatter, splenocytes from SPZ-primed class II<sup>-/-</sup> donors proliferate within 48h. exposure to PfGPI (unshaded) compared with medium controls (shaded). The responding cells are NK1.1<sup>+</sup>, CD4<sup>+</sup>, and include a V $\alpha$ 14<sup>+</sup>, CD4<sup>+</sup> subset. (B) Splenocytes from class II<sup>-/-</sup> donors were exposed to various antigens and [<sup>3</sup>H]TdR incorporation determined after 4 days. Other cultures were exposed to PfGPI or Thy-1 GPI for 4 days, washed and cultured in 10U/ml IL-2 for 2 days, followed by replating with irradiated wild-type APCs and restimulation with either PfGPI or Thy-1 GPI for 48 hours. IL-4 levels in the supernatant were determined by capture ELISA.

**Figure 5.** The proliferative and IL-4 response of NKT cells to PfGPI is MHC-independent and CD1-restricted. (A)  $2 \times 10^4$  sorted NK1.1<sup>+</sup>, CD4<sup>+</sup> cells from wild-type or class II<sup>-/-</sup> donors were placed in triplicate with or without purified GPI on irradiated splenocyte APCs from wild-type (WT), class II<sup>-/-</sup>, b<sub>2</sub>M<sup>-/-</sup> or CD1<sup>-/-</sup> donors, or CD1.1-transfected and sham-transfected J774 macrophages. Tritiated thymidine ([<sup>3</sup>H]TdR) incorporation was determined after 3 days, or IL-4 production in the presence or absence of anti-CD1 was determined as above. (B) Splenocytes from SPZ-primed Class II<sup>-/-</sup> donors were exposed to PfGPI in the presence or absence of anti-CD1 or isotype control and taken for FACS analysis after 3 days.

10

**Figure 6.** CD1-restricted antibody formation to neo-GPI-proteins and malaria SPZ. (A) Donor *nu/nu* mice were primed twice with *P. berghei* SPZ or twice with LPS<sup>FLU</sup>. Splenocytes were cultured in the presence of 10U/ml IL-2, with and without antigen (0.1mg/ml sham-OVA<sup>FLU</sup>, PfGPI-OVA<sup>FLU</sup>, or  $5 \times 10^4$  SPZ), anti-Class I, anti-Class II and anti-CD1, with  $10^4$  NKT cells from SPZ-primed Class II<sup>-/-</sup> donors. Antigen-specific IgG production was quantified by ELISPOT against fluoresceinated Dog serum albumin for responses to OVA<sup>FLU</sup>, and rCS for responses to SPZ. (B) Responses of CD1.1/CD1.2<sup>-/-</sup> (open circles) and Balb/c wild-type mice (black circles) to SPZ, recCS and mfVSG<sup>FLU</sup>. IgG titers were determined as described.

09830451.073101  
FOI E20 T540E360

**DETAILED DESCRIPTION OF THE INVENTION**

The present invention is predicated, in part, on the surprising determination that the helper T cell response to several parasitic proteins is predominantly regulated through CD1-  
5 restricted recognition of the GPI moiety by CD4+ T cells. This determination has facilitated the development of methodology for application, inter alia, in improving immune responses via regulation of helper T cell stimulation, for example, where the immune response is desired for the purpose of therapeutic or prophylactic vaccination.

10 Accordingly, one aspect of the present invention contemplates a method of activating T cells, said method comprising administering a T cell activating effective amount of a molecule or a complex comprising said molecule which molecule or molecule complex is capable of interacting with CD1 on an immune cell to form an association with CD1 which association activates helper T cells.

15

Preferably, the present invention contemplates a method of activating T cells said method comprising administering a T cell activating effective amount of GPI or derivative or equivalent thereof or a complex comprising GPI or derivative or equivalent thereof which GPI or GPI-complex is capable of interacting with CD1 on an immune cell to form an  
20 association with CD1 which association activates helper T cells.

Reference hereinafter to "activating" helper T cells is a reference to upregulating one or more of the functions which said T cells are capable of performing upon stimulation by an antigen, such as, but not limited to, one or more of cell division, differentiation, cell  
25 surface molecule expression or cytokine production.

Reference to "association" should be understood in its broadest sense to include any form of interaction between a molecule and CD1. Said molecule and CD1 may interact via, for example, a covalent bond, ionic bond, hydrogen bond, van Der Waals forces or other  
30 interactive bonding mechanism.

09830451-073101  
TOTAL " 15402860

- 15 -

To the extent that it is not otherwise specified, reference to "GPI" should be read as including reference to all forms of GPI and derivatives or equivalents thereof. Reference to "derivatives" or "equivalents" should be understood to include reference to fragments, parts, portions, chemical equivalents, mutants, homologs and analogs. Chemical equivalents of GPI can act as a functional analog of GPI. Equivalents may not necessarily be derived from GPI but may share certain conformational similarities. Alternatively chemical equivalents may be specifically designed to mimic certain physiochemical properties of GPI. Chemical equivalents may be chemically synthesised or may be detected following, for example, natural product screening. Equivalents also include synthetic carbohydrates and peptide mimetics. Homologs of GPI contemplated herein include, but are not limited to, GPI from different species.

GPI molecules suitable for use in the present invention may be derived from any natural or synthetic source. This includes, for example, GPI moieties derived by genetic manipulation of expression systems and by manipulations of the GPI post-translational modification of proteins via recombinant DNA techniques such as glycosylation inhibitors. Examples of GPI moieties suitable for use in the present invention include but are not limited to microorganism GPI moieties which cause disease conditions such as, the parasitic, fungal and yeast taxa *Plasmodium*, *Trypanosoma*, *Leishmania*, *Toxoplasma* and *Candida*. Preferably, said GPI is a parasite GPI and even more preferably a *Plasmodium* GPI.

According to this preferred embodiment, the present invention contemplates a method of activating helper T cells said method comprising administering a T cell activating effective amount of *Plasmodium* GPI or derivative or equivalent thereof or a complex comprising said *Plasmodium* GPI or derivative or equivalent thereof which *Plasmodium* GPI or *Plasmodium* GPI complex is capable of interacting with CD1 on an immune cell to form an association with CD1 which association activates helper T cells.

Most preferably said *Plasmodium* is *P. falciparum*.

- 16 -

In another most preferred embodiment, said GPI comprises a structure selected from:

- EtN-P-[M $\alpha$ 2]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y  
 EtN-P-[M $\alpha$ 2][G]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y  
 5 EtN-P-[M $\alpha$ 2][X]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y  
 EtN-P-[M $\alpha$ 2][EtN-P]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine  
 10 including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

15

In another preferred embodiment, said GPI comprises a structure selected from:

- EtN-P-M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y  
 M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y  
 20 EtN-P-M $\alpha$ 2M $\alpha$ 6M-Y

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine  
 25 including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

30 In still another preferred embodiment, said GPI comprises a structure selected from:



- 17 -

EtN-P-[M $\alpha$ 2][G]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-YEtN-P-[M $\alpha$ 2][X]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-YEtN-P-[M $\alpha$ 2][EtN-P]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y

5 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 10 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In yet another preferred embodiment, said GPI comprises a structure selected from:

15 M $\alpha$ 2[M $\alpha$ 2][G]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y  
 M $\alpha$ 2[M $\alpha$ 2][X]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y  
 M $\alpha$ 2[M $\alpha$ 2][EtN-P]M $\alpha$ 6M $\alpha$ 4G-Y

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is  
 20 mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y  
 25 is any lipid or phospholipid.

In still yet another preferred embodiment, said GPI comprises a structure selected from:

M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y  
 30 M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y

09830451.073101

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 5 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In a further embodiment, said GPI comprises a structure selected from:


10



15 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 20 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In another further preferred embodiment, said GPI comprises a structure selected from:

25




or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is  
 30 mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or

inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

5


In still another further preferred embodiment, said GPI comprises a structure selected from:



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine  
15 including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

20

In still yet another further preferred embodiment, said GPI comprises a structure selected from:



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
30 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional

- 20 -

linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In another preferred embodiment, said GPI comprises a structure selected from:

5

EtN-P-[M $\alpha$ 2][G]M $\alpha$ 2M-Y

EtN-P-[M $\alpha$ 2][X]M $\alpha$ 2M-Y

EtN-P-[M $\alpha$ 2][EtN-P]M $\alpha$ 2M-Y

10 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
15 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

Reference to "lipid" or "phospholipid" should be understood in its broadest sense and includes, but is not limited to, diacylglycerol, alkylacylglycerol, monoalkylglycerol,  
20 ceramide, sphingolipids and phospholipids such as phosphatidylethanolamine, phosphatidylcholine and phosphatidylserine.

It should also be understood any of these preferred structures may be further modified by substituents of positive, negative or neutral charge such as phosphates, phosphoglycerol,  
25 hexosamines, amino acids, thiols etc in any position and with any type of linkage. This may be particularly useful in the generation of self GPI structures which are typically more highly substituted along the length of the glycan than protozoal GPIs.

GPI "complex" is a reference to a GPI moiety coupled to any other molecule. Said  
30 molecule may be any molecule to which an immune response is sought, for example, a carbohydrate or a peptide, polypeptide or protein such as, but not limited to, peptides,

09030451 073101

- 21 -

polypeptides or proteins naturally anchored to GPI moieties (for example, malarial CS protein, MSP-1, MSP-2, *Leishmanial* PSA-2 or GP63) or any peptides, polypeptides or proteins artificially coupled to a GPI moiety (for example an influenza antigen). Said molecule and said GPI moiety may be covalently linked or may be linked by ionic, hydrogen or other interactive bonding mechanisms. Coupling may be achieved by a variety of techniques including, but in no way limited to, use of a specific expression system or via chemical synthesis. Preferably, said molecule is a protein.

Reference to an "immune cell" should be understood as a reference to any cell of the immune system such as, but not limited to, myeloid cells, stromal cells or antigen presenting cells (for example macrophages).

Reference to "helper T cells" should be understood as a reference to any cell expressing a T cell receptor (expression of a "T cell receptor" is defined as the expression of one or more of an  $\alpha$ ,  $\beta$ ,  $\gamma$  and/or  $\delta$  T cell receptor chain in either homodimeric or heterodimeric form) which can become activated via a CD1-restricted recognition pathway instead of, or in addition to, a capacity to become activated via a MHC II restricted recognition pathway and which acts to stimulate, upregulate or otherwise modulate any aspect of the immune response via any one or more of a variety of mechanisms including, for example, cell-cell contact or production of soluble mediators. Said T cells include, but are not limited to, thymically derived T cells. Preferably said T cells express CD4 and even more preferably CD4 and NK1.1. This cell type is often referred to in the art as the "NKT cell".

According to this most preferred embodiment, the present invention contemplates a method of activating helper T cells said method comprising administering a T cell activating effective amount of GPI or derivative or equivalent thereof or a complex comprising said GPI or derivative or equivalent thereof which GPI or GPI complex is capable of interacting with CD1 on an immune cell to form an association with CD1 which association activates CD4+ NK1.1+ T cells.

30

Without limiting the present invention to any one theory or mode of action, administration

- 22 -

of GPI or a GPI complex leads to presentation of the GPI moiety, or a molecule complexed to the GPI moiety, by a non-polymorphic CD1 restriction element. CD1 is expressed on a variety of cells including, for example, macrophages. Coupling of the GPI moiety (or the molecule complexed to the GPI) to the CD1 may be by covalent bonding. Recognition of the CD1-GPI unit by a subclass of CD4<sup>+</sup> T cells leads to their activation. Said T cells do not recognise the GPI moiety or the molecule complexed to the GPI via the traditional MHC II restricted route of presentation, instead said T cells are activated through CD1-restricted recognition. Said T cells represent a subset of the T cell population of an individual. For example, T cells exhibiting the phenotype CD4<sup>+</sup>, NK1.1<sup>+</sup> are able to become activated via the CD1-restricted recognition pathway. The present invention should be understood to extend to methods of activating T cells by administration of GPI or GPI complex wherein said T cells are activated by one or both of CD1 or MHC II restricted recognition.

Even more preferably the present invention contemplates a method of activating CD4<sup>+</sup>, NK1.1<sup>+</sup> T cells said method comprising administering a T cell activating effective amount of a *Plasmodium* GPI or derivative or equivalent thereof or a complex comprising said *Plasmodium* GPI or derivative or equivalent thereof which *Plasmodium* GPI or *Plasmodium* GPI complex is capable of interacting with CD1 on an immune cell to form an association with CD1 which association activates CD4<sup>+</sup> NK1.1<sup>+</sup> T cells.

Most preferably said *Plasmodium* is *P. falciparum*.

Still without limiting the present invention to any one theory or mode of action, the T cells of the present invention can co-operate with B cells to result in CD1-restricted antibody production, the specificity of said antibody being directed to the GPI moiety (where said GPI is administered in isolation) or to a molecule complexed to the GPI moiety (where a GPI complex is administered). For example, said complexed molecule may be a protein to which an antibody response is desired, such as an influenza antigen. Activation of said B cells is supported by the production of cytokines, such as IL-4, by the CD1-restricted activated T cells. GPI-anchoring, therefore, permits antibody formation by a non-MHC

restricted immunological process. Unlike MHC, which is a highly polymorphic molecule, CD1 is non-polymorphic thereby resulting in little observed non-responsiveness in human populations.

- 5 Accordingly, the present invention contemplates a method of activating T helper cells said method comprising administering a T cell activating effective amount of a GPI or derivative or equivalent thereof or a complex comprising said GPI or derivative or equivalent thereof which GPI or GPI complex is capable of interacting with CD1 on an immune cell to form an association with CD1 which association activates T helper cells  
10 wherein said activated T cells provide B cell help.

Preferably said T helper cells are CD4<sup>+</sup>, NK1.1<sup>+</sup> T cells.

- More preferably, said GPI molecule is a parasitic, fungal or yeast GPI moiety. Most  
15 preferably, said GPI molecule is a parasite GPI and even more preferably a *Plasmodium* GPI.

- Accordingly, a related aspect of the present invention contemplates a method of activating CD4<sup>+</sup>, NK1.1<sup>+</sup> T cells said method comprising administering a T cell activating effective  
20 amount of a *Plasmodium* GPI or derivative or equivalent thereof or a complex comprising said *Plasmodium* GPI or derivative or equivalent thereof which *Plasmodium* GPI or *Plasmodium* GPI complex is capable of interacting with CD1 on an immune cell to form an association with CD1 which association activates CD4<sup>+</sup>, NK1.1<sup>+</sup> T cells wherein said activated T cells provide B cell help.


25

Most preferably said *Plasmodium* is *P. falciparum*.

In another most preferred embodiment, said GPI comprises a structure selected from:


- 30 EtN-P-[M $\alpha$ 2]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y  
EtN-P-[M $\alpha$ 2][G]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y

- 24 -



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is  
 5 mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine  
 including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 linkages which may be substituted with any other positional linkages as required, and Y  
 10 is any lipid or phospholipid.

In another preferred embodiment, said GPI comprises a structure selected from:



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is  
 mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine  
 20 including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 linkages which may be substituted with any other positional linkages as required, and Y  
 is any lipid or phospholipid.

25

In still another preferred embodiment, said GPI comprises a structure selected from:



05830451-073101



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 5 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In yet another preferred embodiment, said GPI comprises a structure selected from:

10



15 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 20 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In still yet another preferred embodiment, said GPI comprises a structure selected from:

25



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine  
 30 including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may

- 26 -

be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

5 In a further embodiment, said GPI comprises a structure selected from:

$M\alpha_2[M\alpha_2]M\alpha_6M\alpha_4G\alpha_6Ino-Y$

$M\alpha_2[M\alpha_2][G]M\alpha_6M\alpha_4G\alpha_6Ino-Y$

$M\alpha_2[M\alpha_2][X]M\alpha_6M\alpha_4G\alpha_6Ino-Y$

10

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 15 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In another further preferred embodiment, said GPI comprises a structure selected from:

20

$EtN-P-[M\alpha_2][G]M\alpha_2M\alpha_6M-Y$

$EtN-P-[M\alpha_2][X]M\alpha_2M\alpha_6M-Y$


$EtN-P-[M\alpha_2][EtN-P]M\alpha_2M\alpha_6M-Y$

25 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 30 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

09630451.073101

- 27 -


In still another further preferred embodiment, said GPI comprises a structure selected from:



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine  
10 including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

15

In still yet another further preferred embodiment, said GPI comprises a structure selected from:



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
25 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

30 In another preferred embodiment, said GPI comprises a structure selected from:

09830451-073101

- 28 -

EtN-P-[M $\alpha$ 2][G]M $\alpha$ 2M-YEtN-P-[M $\alpha$ 2][X]M $\alpha$ 2M-YEtN-P-[M $\alpha$ 2][EtN-P]M $\alpha$ 2M-Y

5 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 10 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

Reference to "B cell help" should be understood as a reference to the provision, by said activated T cells, of signals which act to stimulate, up-regulate or otherwise modulate or  
 15 maintain B cell and/or plasma cell viability or functional activity. Said signals may take any form including, for example, cell/cell contact or the production of soluble mediators such as cytokines.

In addition to co-operating with B cells to support antibody production, the T cells  
 20 activated in accordance with the method of the present invention produce a range of cytokines which can modulate aspects of the immune response other than antibody production. For example, and without limiting the present invention in any way, CD-1 restricted NKT cells produce very high levels of the cytokines IL-4 and IFN- $\gamma$  suggesting that these cells are involved in the downstream regulation of TH1/TH2 differentiation.  
 25 Accordingly, these cells are thought to be involved in the aetiology of those diseases which show a pronounced TH1/TH2 dependence, including cerebral malaria, tuberculosis, leprosy, *leishmaniasis*, type I diabetes, autoimmune arthritis and systemic lupus erythematosus. They are also thought to be involved in tumour rejection.

30 The up-regulation of NKT cell proliferation and differentiation therefore provides a mechanism for skewing the T helper cell response towards a TH1 or a TH2 response.

09830451 073101

Skewing of the CD1 restricted T helper cell response, for example, may be achieved by techniques which are known to those skilled in the art. These techniques include, for example, the route by which the GPI molecule/GPI complex is administered, co-stimulatory signals which are provided together with the GPI molecule/GPI complex and/or the choice of vehicle which is used to administer the GPI molecule/GPI complex. For example, administration of the GPI molecule/GPI complex is alum or liposomes will skew the CD1 restricted T helper response towards a TH2 type response while administration of these molecules utilising ISCOM's will skew the response to either the TH1 or TH2 type response depending on the route of administration which is utilised. The administration of an IL-12 or IFN- $\alpha$  co-stimulatory agent together with the GPI molecule/GPI complex is likely to skew the response towards a TH1 type response while the administration of an IL-4 co-stimulatory agent together with the GPI molecule/GPI complex is more likely to skew the response towards a TH2 type response.

The method of the present invention therefore provides not only a mechanism of activating T helper cells in a non-MHC restricted fashion in response to a GPI stimulus, but the method may also be adapted to skew the nature of the immune response which is ultimately effected. For example, up-regulation of the subject T cells can be designed to provide B cell help or to otherwise effect a TH1 or TH2 type response.

20

Accordingly, another related aspect of the present invention contemplates a method of activating T helper cells said method comprising administering a T cell activating effective amount of a GPI or derivative or equivalent thereof or a complex comprising said GPI or derivative or equivalent thereof which GPI or GPI complex is capable of interacting with CD1 on an immune cell to form an association with CD1 which association activates T helper cells wherein said activated T cells induce or otherwise upregulate a TH1-type response.

Preferably said T helper cell is a CD4<sup>+</sup> NK 1.1<sup>+</sup> cell. More preferably said GPI molecule is a parasitic, fungal or yeast GPI moiety and even more preferably a *Plasmodium* GPI.

- 30 -



Yet another related aspect of the present invention contemplates a method of activating T helper cells said method comprising administering a T cell activating effective amount of a GPI or derivative or equivalent thereof or a complex comprising said GPI or derivative or equivalent thereof which GPI or GPI complex is capable of interacting with CD1 on an immune cell to form an association with CD1 which association activates T helper cells wherein said activated T cells induce or otherwise upregulate a TH2-type response.

Preferably said T helper cell is a CD4<sup>+</sup> NK1.1<sup>+</sup> cell. More preferably said GPI molecule is a parasitic, fungal or yeast GPI moiety and even more preferably a *Plasmodium* GPI.

10

Reference to a "TH1-type" or "TH2-type" response should be understood as a reference to a T helper cell response which exhibits one or more of the functional characteristics of a TH1 or TH2 response, respectively, and which is directly or indirectly mediated by the CD1 restricted T cells which are activated in accordance with the method of the present invention. These characteristics include, but are not limited to the cytokine profile which is induced in the subject. For example, a TH1 response is characterised by the up-regulation of IFN- $\gamma$  expression while a TH2 response is characterised by the up-regulation of IL-4 production.

20 In a most preferred embodiment, said GPI comprises a structure selected from:



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional

09830451.073101

linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In another preferred embodiment, said GPI comprises a structure selected from:

5

EtN-P-M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y

M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y

EtN-P-M $\alpha$ 2M $\alpha$ 6M-Y

10 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
15 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In still another preferred embodiment, said GPI comprises a structure selected from:

20 EtN-P-[M $\alpha$ 2][G]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y  
EtN-P-[M $\alpha$ 2][X]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y  
EtN-P-[M $\alpha$ 2][EtN-P]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is  
25 mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y  
30 is any lipid or phospholipid.

05830451.073101  
T01E20.T540E860


- 32 -

In yet another preferred embodiment, said GPI comprises a structure selected from:




or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
 10 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

15 In still yet another preferred embodiment, said GPI comprises a structure selected from:



20 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 25 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In a further embodiment, said GPI comprises a structure selected from:



09830451.073101





or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

10

In another further preferred embodiment, said GPI comprises a structure selected from:



15



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

25 In still another further preferred embodiment, said GPI comprises a structure selected from:



30



09830451.073101

- 34 -

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 5 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In still yet another further preferred embodiment, said GPI comprises a structure selected  
 10 from:

$$M\alpha_2M\alpha_6M-Y$$

$$M\alpha_6M\alpha_4G-Y$$

15 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 20 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In another preferred embodiment, said GPI comprises a structure selected from:

$$25 \quad EtN-P-[M\alpha_2][G]M\alpha_2M-Y$$

$$EtN-P-[M\alpha_2][X]M\alpha_2M-Y$$

$$EtN-P-[M\alpha_2][EtN-P]M\alpha_2M-Y$$

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is  
 30 mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or

09830451.073101

inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

5

A further aspect of the present invention relates to the use of the invention in relation to disease conditions. For example:

- 10 (i) As therapeutic or prophylactic immunogenic compositions directed to molecules naturally GPI-anchored such as, for example, the malarial CS protein, MSP-I, and MSP-2, *Leishmanial* PSA-2 and gp63.
- (ii) As therapeutic or prophylactic immunogenic compositions directed to GPI molecules themselves, for example in treating parasitic, fungal and yeast infections.
- 15 (iii) To enable antibody production where MHC II restricted T cell activation is undesirable, such as where said MHC II restricted T cell would engender an autoimmune reaction. For example, the M protein of Group A Streptococcus exhibits T cell epitopes which are cross-reactive with human heart tissue but
- 20 exhibits B cell epitopes protective against the bacterium. Said B cell epitope may be coupled to GPI to facilitate non-MHC restricted antibody formation.
- (iv) To provide a universal T cell epitope for any therapeutic or prophylactic immunogenic compositions construct.

25

In one example, the use of the present invention to regulate a TH1/TH2 response provides a mechanism to therapeutically or prophylactically treat disease conditions which show a pronounced TH1/TH2 dependence such as, but not limited to, cerebral malaria, tuberculosis, leprosy, *leishmaniasis*, type I diabetes, autoimmune arthritis, systemic lupus

30 erythromatosis. In particular, skewing towards or inducing a TH1 response in subjects suffering from neoplasia, cancer or *Leishmaniasis* is desirable. Skewing towards or

- 36 -

induction of a TH2 response is desirable where a subject is suffering from SLE, type I diabetes, autoimmune arthritis and cerebral malaria, for example. The method of the present invention may also be utilised to provide B cell help, thereby leading to an antibody response which is useful, for example, in the treatment or prophylaxis of parasitic  
5 infections such as malaria and *leishmaniasis*. The production of B cell help can be achieved, for example, by inducing a TH2 response.

Accordingly, in another aspect, the present invention provides a method of inducing, in a mammal, an immune response directed to a GPI said method comprising administering  
10 to said mammal a T cell activating effective amount of GPI or derivative or equivalent thereof which GPI is capable of interacting with CD1 on an immune cell to form an association with CD1 which association activates helper T cells.

Preferably, said T cell is a CD4+ T cell and even more preferably a CD4+ NK1.1+ T  
15 cell. In another preferred embodiment, said activated T cell provides B cell help.

Still more preferably said GPI is *Plasmodium* GPI.

In still another aspect, the present invention provides a method of inducing, in a mammal,  
20 an immune response directed to an antigen, said method comprising administering to said mammal a T cell activating effective amount of GPI or derivative or equivalent thereof complexed to said antigen, which GPI-antigen complex is capable of interacting with CD1 on an immune cell to form an association with CD1, which association activates helper T cells.

25

Preferably, said T cell is a CD4+ T cell and even more preferably a CD4+ NK1.1+ T cell.

In another preferred embodiment, said activated T cell provides B cell help.

30

In still another preferred embodiment, said activated T cell induces or otherwise up-

09330451-073101  
TOTAL 20 75406860

- 37 -

regulates a TH1 or TH2 response.

In another most preferred embodiment, said GPI comprises a structure selected from:

- 5        EtN-P-[M $\alpha$ 2]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y  
        EtN-P-[M $\alpha$ 2][G]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y  
        EtN-P-[M $\alpha$ 2][X]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y  
        EtN-P-[M $\alpha$ 2][EtN-P]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y

10 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is  
 mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine  
 including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 15 linkages which may be substituted with any other positional linkages as required, and Y  
 is any lipid or phospholipid.

In another preferred embodiment, said GPI comprises a structure selected from:

- 20        EtN-P-M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y  
        M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y  
        EtN-P-M $\alpha$ 2M $\alpha$ 6M-Y

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is  
 25 mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine  
 including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 linkages which may be substituted with any other positional linkages as required, and Y  
 30 is any lipid or phospholipid.

09830451.073101

- 38 -

In still another preferred embodiment, said GPI comprises a structure selected from:



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
10 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

15 In yet another preferred embodiment, said GPI comprises a structure selected from:



20

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
25 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In still yet another preferred embodiment, said GPI comprises a structure selected from:

30



09830451.073101

- 39 -



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

10

In a further embodiment, said GPI comprises a structure selected from:



15



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

25 In another further preferred embodiment, said GPI comprises a structure selected from:



30

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is

09830451.073101

- 40 -

mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In still another further preferred embodiment, said GPI comprises a structure selected from:

10



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In still yet another further preferred embodiment, said GPI comprises a structure selected from:

25



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or



09330451.073101



inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

5

In another preferred embodiment, said GPI comprises a structure selected from:



10



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
 15 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

20 Accordingly, another aspect of the present invention contemplates a method of treating a mammal said method comprising administering to said mammal an effective amount of GPI or derivative or equivalent thereof or a complex comprising said GPI or derivative or equivalent thereof which GPI or GPI complex is capable of interacting with CD1 on an immune cell to form an association with CD1 which association activates helper T cells.

25

In one preferred embodiment said mammal has a parasitic infection and more preferably a *Plasmodium* infection. Most preferably said activated helper T cells provide B cell help.

Still more preferably said GPI is *Plasmodium* GPI.

30

In another preferred embodiment, said mammal is in need of the induction, modulation or

09830451.073101

- 42 -

skewing of a helper T cell response along the TH1 or TH2 pathway and said activated T helper cells induce, modulate or skew the T helper response along the TH1 or TH2 pathway.

- 5 In another aspect the present invention contemplates a method of treating a mammal said method comprising administering a mammalian helper T cell activation effective amount of a GPI or derivative or equivalent thereof or a complex comprising said GPI or derivative or equivalent thereof which GPI or GPI complex is capable of interacting with CD1 on an immune cell to form an association with CD1 which association activates helper T cells.

10

In one preferred embodiment said mammal has a parasitic infection and more preferably a *Plasmodium* infection. Most preferably said activated helper T cells provide B cell help.

Still more preferably said GPI is *Plasmodium* GPI.

15

In another preferred embodiment, said mammal is in need of the induction, modulation or skewing of a helper T cell response along the TH1 or TH2 pathway and said activated T helper cells induce, modulate or skew the T helper response along the TH1 or TH2 pathway.

20

Yet another aspect of the present invention contemplates the use of GPI or derivative or equivalent thereof or a complex comprising said GPI or derivative or equivalent thereof which GPI or GPI complex is capable of interacting with CD1 on an immune cell, in the manufacture of a medicament for the activation of helper T cells in a mammal.

25

In one preferred embodiment said mammal has a parasitic infection and more preferably a *Plasmodium* infection. Most preferably said activated helper T cells provide B cell help.

Still more preferably said GPI is *Plasmodium* GPI.

30

In another preferred embodiment, said mammal is in need of the induction, modulation or

09830451-073101  
TOTAL 15408860

- 43 -

skewing of a helper T cell response along the TH1 or TH2 pathway and said activated T helper cells induce, modulate or skew the T helper response along the TH1 or TH2 pathway.

- 5 The term "mammal" includes humans, primates, livestock animals (e.g. horses, cattle, sheep, pigs and donkeys) laboratory test animals (e.g. mice, rats, rabbits, guinea pigs) companion animals (e.g. dogs and cats) and captive wild animals (e.g. kangaroos, deer, foxes). Preferably, the mammal is a human or laboratory test animal. Even more preferably the mammal is a human.

10

In a related aspect of the present invention the mammal undergoing treatment may be human or an animal in need of therapeutic or prophylactic treatment.

- Accordingly, the present invention provides a method for the treatment and/or prophylaxis  
15 of a mammalian disease condition characterised by a micro-organism infection, said method comprising administering to said mammal an effective amount of GPI or derivative or equivalent thereof or a complex comprising said GPI or derivative or equivalent thereof which GPI or GPI complex is capable of interacting with CD1 on an immune cell to form an association with CD1 which association activates helper T cells.

20

Preferably said microorganism infection is a parasitic infection and more preferably a *Plasmodium* infection. Still more preferably said activated T cells provide B cell help.

- In another aspect, the present invention provides a method for the treatment and/or  
25 prophylaxis of a mammalian disease condition characterised by the insufficiency or absence of an appropriate TH1 response said method comprising administering to said mammal an effective amount of GPI or derivative or equivalent thereof or a complex comprising said GPI or derivative or equivalent thereof which GPI or GPI complex is capable of interacting with CD1 on an immune cell to form an association with CD1 which association induces  
30 or otherwise upregulates a TH1 response.

09830451.073101  
TOTAL 754360

- 44 -

In still another aspect, the present invention provides a method for the treatment and/or prophylaxis of a mammalian disease condition characterised by the insufficiency or absence of an appropriate TH2 response said method comprising administering to said mammal an effective amount of GPI or derivative or equivalent thereof or a complex comprising said GPI or derivative or equivalent thereof which GPI or GPI complex is capable of interacting with CD1 on an immune cell to form an association with CD1 which association induces or otherwise upregulates a TH2 response.

Reference to a disease condition "characterised by the insufficiency or absence of an appropriate" TH1 or TH2 response should be understood as a reference to a disease condition in which the appropriate response is either not functional or else is functional at a level too low to be therapeutically or prophylactically effective. It should also be understood as a reference to a condition where one form of helper T cell response is in effect but where it is desirable to skew the response to the other form (for example, skewing a TH1 response to a TH2 response or vice versa). The skewing of the helper T cell response is particularly useful in disease conditions where one or more of the symptoms exhibited by a patient are due to the functional features characteristic of a TH1 or TH2 response, for example the cytokine profile.

An "effective amount" means an amount necessary at least partly to attain the desired immune response, or to prevent or to delay the onset or inhibit progression or halt altogether, the onset or progression of a particular condition being treated. This amount varies depending upon the health and physical condition of the individual to be treated, the taxonomic group of individual to be treated, the capacity of the individual's immune system to synthesise antibodies, the degree of protection desired, the formulation of the vaccine, the assessment of the medical situation, and other relevant factors. It is expected that the amount will fall in a relatively broad range that can be determined through routine trials.

Reference herein to "treatment" and "prophylaxis" is to be considered in its broadest context. The term "treatment" does not necessarily imply that a mammal is treated until

- 45 -

total recovery. Similarly, "prophylaxis" does not necessarily mean that the subject will not eventually contract a disease condition. Accordingly, treatment and prophylaxis include amelioration of the symptoms of a particular condition or preventing or otherwise reducing the risk of developing a particular condition. The term "prophylaxis" may be considered  
 5 as reducing the severity of onset of a particular condition. "Treatment" may also reduce the severity of an existing condition or the frequency of acute attacks (for example, reducing the severity of initial infection).

In accordance with these methods, the GPI or GPI complex defined in accordance with the  
 10 present invention may be coadministered with one or more other compounds or molecules. By "coadministered" is meant simultaneous administration in the same formulation or in two different formulations via the same or different routes or sequential administration by the same or different routes. By "sequential" administration is meant a time difference of from seconds, minutes, hours or days between the administration of the two types of  
 15 molecules. These molecules may be administered in any order.

The present invention should also be understood to extend to immunogenic compositions for use in the methods as hereinbefore defined.

20 Accordingly, in a related aspect, the present invention is directed to a composition which activates T cells, said composition comprising a GPI or derivative or equivalent thereof or a complex comprising GPI or derivative or equivalent thereof which GPI or GPI-complex is capable of interacting with CD1 on an immune cell to form an association with CD1 which association activates helper T cells.

25


Preferably said GPI is *Plasmodium* GPI.

In another most preferred embodiment, said GPI comprises a structure selected from:

30 EtN-P-[M $\alpha$ 2]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y  
 EtN-P-[M $\alpha$ 2][G]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y


09330451.073101

- 46 -



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is  
 5 mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine  
 including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 linkages which may be substituted with any other positional linkages as required, and Y  
 10 is any lipid or phospholipid.

In another preferred embodiment, said GPI comprises a structure selected from:



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is  
 mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine  
 20 including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 linkages which may be substituted with any other positional linkages as required, and Y  
 is any lipid or phospholipid.

25

In still another preferred embodiment, said GPI comprises a structure selected from:



09330451-073101

- 47 -

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In yet another preferred embodiment, said GPI comprises a structure selected from:

10



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In still yet another preferred embodiment, said GPI comprises a structure selected from:

25



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may

09830451.073101

be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

5 In a further embodiment, said GPI comprises a structure selected from:

$M\alpha_2[M\alpha_2]M\alpha_6M\alpha_4G\alpha_6Ino-Y$

$M\alpha_2[M\alpha_2][G]M\alpha_6M\alpha_4G\alpha_6Ino-Y$

$M\alpha_2[M\alpha_2][X]M\alpha_6M\alpha_4G\alpha_6Ino-Y$

10

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 15 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In another further preferred embodiment, said GPI comprises a structure selected from:

20

$EtN-P-[M\alpha_2][G]M\alpha_2M\alpha_6M-Y$

$EtN-P-[M\alpha_2][X]M\alpha_2M\alpha_6M-Y$


$EtN-P-[M\alpha_2][EtN-P]M\alpha_2M\alpha_6M-Y$

25 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 30 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.



- 49 -


In still another further preferred embodiment, said GPI comprises a structure selected from:



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

15

In still yet another further preferred embodiment, said GPI comprises a structure selected from:



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

30 In another preferred embodiment, said GPI comprises a structure selected from:

09830451.073101

- 50 -

EtN-P-[M $\alpha$ 2][G]M $\alpha$ 2M-YEtN-P-[M $\alpha$ 2][X]M $\alpha$ 2M-YEtN-P-[M $\alpha$ 2][EtN-P]M $\alpha$ 2M-Y

5 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 10 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In another aspect, the present invention relates to an immunogenic composition comprising as the active component GPI or derivative or equivalent thereof or GPI complex, as  
 15 broadly described above.

Preferably, said GPI is *Plasmodium* GPI.

In another most preferred embodiment, said GPI comprises a structure selected from:  
 20

EtN-P-[M $\alpha$ 2]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-YEtN-P-[M $\alpha$ 2][G]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-YEtN-P-[M $\alpha$ 2][X]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-YEtN-P-[M $\alpha$ 2][EtN-P]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y

25

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 30 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y

09030451.073101

is any lipid or phospholipid.

In another preferred embodiment, said GPI comprises a structure selected from:

- 5        EtN-P-M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y  
          M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y  
          EtN-P-M $\alpha$ 2M $\alpha$ 6M-Y

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is  
 10 mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine  
 including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 linkages which may be substituted with any other positional linkages as required, and Y  
 15 is any lipid or phospholipid.

In still another preferred embodiment, said GPI comprises a structure selected from:

- EtN-P-[M $\alpha$ 2][G]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y  
 20        EtN-P-[M $\alpha$ 2][X]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y  
          EtN-P-[M $\alpha$ 2][EtN-P]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is  
 mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine  
 25 including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 linkages which may be substituted with any other positional linkages as required, and Y  
 is any lipid or phospholipid.

30

In yet another preferred embodiment, said GPI comprises a structure selected from:

09830451.073101  
 101E2015409860

- 52 -

$$M\alpha_2[M\alpha_2][G]M\alpha_2M\alpha_6M\alpha_4G-Y$$

$$M\alpha_2[M\alpha_2][X]M\alpha_2M\alpha_6M\alpha_4G-Y$$

$$M\alpha_2[M\alpha_2][EtN-P]M\alpha_6M\alpha_4G-Y$$

5 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
10 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In still yet another preferred embodiment, said GPI comprises a structure selected from:

15  $M\alpha_6M\alpha_4G\alpha_6Ino-Y$   
 $M\alpha_2M\alpha_6M\alpha_4G\alpha_6Ino-Y$

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine  
20 including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

25

In a further embodiment, said GPI comprises a structure selected from:

$M\alpha_2[M\alpha_2]M\alpha_6M\alpha_4G\alpha_6Ino-Y$   
 $M\alpha_2[M\alpha_2][G]M\alpha_6M\alpha_4G\alpha_6Ino-Y$   
 30  $M\alpha_2[M\alpha_2][X]M\alpha_6M\alpha_4G\alpha_6Ino-Y$

09830451-073101

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
 5 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In another further preferred embodiment, said GPI comprises a structure selected from:

10



15 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
 20 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In still another further preferred embodiment, said GPI comprises a structure selected from:

25



30 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine

09830451-073101

- 54 -

including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y  
5 is any lipid or phospholipid.

In still yet another further preferred embodiment, said GPI comprises a structure selected from:

10  $M\alpha_2M\alpha_6M-Y$   
 $M\alpha_6M\alpha_4G-Y$

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine  
15 including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

20

In another preferred embodiment, said GPI comprises a structure selected from:

$EtN-P-[M\alpha_2][G]M\alpha_2M-Y$   
 $EtN-P-[M\alpha_2][X]M\alpha_2M-Y$   
 25  $EtN-P-[M\alpha_2][EtN-P]M\alpha_2M-Y$

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
30 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional

0930451-03101  
 10720" 1540360

- 55 -

linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

Still another aspect of the present invention is directed to a pharmaceutical composition capable of activating T cells, said composition comprising a GPI or derivative or equivalent thereof or a complex comprising GPI or derivative or equivalent thereof which GPI or GPI-complex is capable of interacting with CD1 on an immune cell to form an association with CD1, which association activates helper T cells, together with one or more pharmaceutically acceptable carriers and/or diluents.

10

Preferably, said GPI is *Plasmodium* GPI.

In another most preferred embodiment, said GPI comprises a structure selected from:

15

EtN-P-[M $\alpha$ 2]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y

EtN-P-[M $\alpha$ 2][G]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y

EtN-P-[M $\alpha$ 2][X]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y

EtN-P-[M $\alpha$ 2][EtN-P]M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G $\alpha$ 6Ino-Y

20 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
25 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In another preferred embodiment, said GPI comprises a structure selected from:

30

EtN-P-M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y

M $\alpha$ 2M $\alpha$ 6M $\alpha$ 4G-Y

09630451.073101

- 56 -

EtN-P-M $\alpha$ 2M $\alpha$ 6M-Y

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

10

In still another preferred embodiment, said GPI comprises a structure selected from:



15



or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

25 In yet another preferred embodiment, said GPI comprises a structure selected from:



30

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is

09630451-073101



mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In still yet another preferred embodiment, said GPI comprises a structure selected from:

10  $M\alpha_6M\alpha_4G\alpha_6Ino-Y$

$M\alpha_2M\alpha_6M\alpha_4G\alpha_6Ino-Y$

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

20

In a further embodiment, said GPI comprises a structure selected from:

$M\alpha_2[M\alpha_2]M\alpha_6M\alpha_4G\alpha_6Ino-Y$

$M\alpha_2[M\alpha_2][G]M\alpha_6M\alpha_4G\alpha_6Ino-Y$

25  $M\alpha_2[M\alpha_2][X]M\alpha_6M\alpha_4G\alpha_6Ino-Y$

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional


09830451.073101

- 58 -

linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In another further preferred embodiment, said GPI comprises a structure selected from:

5



10 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
15 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

In still another further preferred embodiment, said GPI comprises a structure selected from:

20



25 or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional  
30 linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

09330451.073101

In still yet another further preferred embodiment, said GPI comprises a structure selected from:

$M\alpha_2M\alpha_6M-Y$

5  $M\alpha_6M\alpha_4G-Y$

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or  
10 inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

15 In another preferred embodiment, said GPI comprises a structure selected from:

$EtN-P-[M\alpha_2][G]M\alpha_2M-Y$

$EtN-P-[M\alpha_2][X]M\alpha_2M-Y$

$EtN-P-[M\alpha_2][EtN-P]M\alpha_2M-Y$

20

or derivative or equivalent thereof wherein EtN is ethanolamine, P is phosphate, M is mannose, G is non-N-acetylated glucosamine, [G] is any non-N-acetylated hexosamine including glucosamine, or any other nitrous-acid labile substituent, Ino is inositol or inositol-phosphoglycerol, [X] is any other substituent,  $\alpha$  represents  $\alpha$ -linkages which may  
25 be substituted with  $\beta$ -linkages wherever required, numeric values represent positional linkages which may be substituted with any other positional linkages as required, and Y is any lipid or phospholipid.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions  
30 (where water soluble) and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion or may be in the form of a cream or other form suitable

- 60 -

- for topical application. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The preventions of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.
- 15 Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilisation. Generally, dispersions are prepared by incorporating the various sterilised active ingredient into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze-drying technique which yield a powder of the active ingredient plus any additional desired ingredient from previously sterile-filtered solution thereof.
- 25 When the active ingredients are suitably protected they may be orally administered, for example, with an inert diluent or with an assimilable edible carrier, or it may be enclosed in hard or soft shell gelatin capsule, or it may be compressed into tablets, or it may be incorporated directly with the food of the diet. For oral therapeutic administration, the active compound may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 1% by weight of active

- 61 -

compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 5 to about 80% of the weight of the unit. The amount of active compound in such therapeutically useful compositions in such that a suitable dosage will be obtained. Preferred compositions or preparations according to the  
5 present invention are prepared so that an oral dosage unit form contains between about 0.1  $\mu$ g and 2000 mg of active compound.

The tablets, troches, pills, capsules and the like may also contain the components as listed hereafter: a binder such as gum, acacia, corn starch or gelatin; excipients such as dicalcium  
10 phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin may be added or a flavouring agent such as peppermint, oil of wintergreen, or cherry flavouring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may  
15 be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the active compound, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavouring such as cherry or orange flavour. Of course, any material used in preparing any dosage unit form should be pharmaceutically  
20 pure and substantially non-toxic in the amounts employed. In addition, the active compound(s) may be incorporated into sustained-release preparations and formulations.

Pharmaceutically acceptable carriers and/or diluents include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption  
25 delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, use thereof in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

30

It is especially advantageous to formulate parenteral compositions in dosage unit form for

- 62 -

ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the novel dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active material and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active material for the treatment of disease in living subjects having a diseased condition in which bodily health is impaired as herein disclosed in detail.

10

The principal active ingredient is compounded for convenient and effective administration in effective amounts with a suitable pharmaceutically acceptable carrier in dosage unit form as hereinbefore disclosed. A unit dosage form can, for example, contain the principal active compound in amounts ranging from 0.5  $\mu\text{g}$  to about 2000 mg. Expressed in proportions, the active compound is generally present in from about 0.5  $\mu\text{g}$  to about 2000 mg/ml of carrier. In the case of compositions containing supplementary active ingredients, the dosages are determined by reference to the usual dose and manner of administration of the said ingredients.

20 The pharmaceutical composition may also comprise genetic molecules such as a vector capable of transfecting target cells where the vector carries a nucleic acid molecule capable of expressing, for example, a functional equivalent to a GPI or derivative thereof. The vector may, for example, be a viral vector and it may be administered by any suitable method including, for example transfection directly into the cells of the mammal being  
25 treated or transfection into a host cell, such as a bacterium, yeast or attenuated parasite, which is then introduced into the mammal.

The present invention is further described by the following non-limiting Examples:

- 63 -

### EXAMPLE 1

#### REAGENTS AND ANIMALS

Pronase was obtained from Boehringer Mannheim. Octyl-Sepharose, Protein-G  
5 Sepharose, *n*-octylthioglucopyranoside (*n*-otg), phenylmethanesulfonyl fluoride (PMSF),  
*p*-tosyl-L-lysinechloromethylketone (TLCK), N-tosyl-L-phenylalaninechloro-  
methylketone (TPCK), *p*-chloromercuriphenylsulphonic acid (*p*-CMPS), aprotinin,  
leupeptin, pepstatin, iodoacetamide, iodoacetic acid and *n*-ethylmaleimide (NEM) were  
obtained from Sigma Chemical Co. Sephadex was from Pharmacia. Analytical or  
10 HPLC grade, acetic acid, butanol, chloroform, diethyl ether, ethanol, methanol and  
water were obtained from BDH and Waters. Silica G60 TLC plates were from Merck  
Darmstadt. Tritiated mannose, glucosamine, myristic and palmitic acids were from  
Amersham. The recombinant *P. falciparum* CS protein 2.3 consists of the entire gene  
except for the C-terminal 21 amino acids. The *P. berghei* rCS encompasses amino  
15 acids 81-277, including the central repetitive domain. The NANP<sub>40</sub> peptide, and the  
17-mer peptide of the tandemly repeating domain of the *P. berghei* CS protein  
(DPPPPNPN)<sub>2</sub>D, were synthesized by routine methods.

Adult female C57Bl/6 wild-type mice, C57Bl/6 lacking the MHC Class II gene, or the  
20 CD1.1 and CD1.2 genes, congenic mice on the C57Bl/10 or Balb background, Balb/c  
*nu/nu* mice and other mouse haplotypes used in the study were maintained in specific  
pathogen free animal facilities.

### EXAMPLE 2

#### CHIMERAS

25 Bone marrow was derived from the femurs and tibiae of Balb/c, Balb/B, Balb/K  
donors. T cells and NK cells were depleted by complement-mediated lysis of Thy-1<sup>+</sup>,  
CD4<sup>+</sup>, Lyt.2<sup>+</sup> and asialo-GM<sup>+</sup> cells. Bone marrow aspirates in RPMI 1640 + 3%  
30 BSA at 10<sup>7</sup> cells/ml were incubated on ice with the appropriate dilution of specific  
antibody, followed by centrifugation at 2°C and resuspension in low toxicity, mouse

- 64 -

lymphocyte absorbed, sterile-filtered Guinea pig complement at 37°C in RPMI 1640 + 3% BSA for 1 hr. Syngeneic or allogeneic recipients, maintained on acidified water and tetracycline, were irradiated by cobalt source (1000 Rads) and inoculated i.v. with 10<sup>7</sup> T-depleted cells. Animals were maintained on acidified water and rested for 12 weeks prior to testing for chimerism and use in experiments. To generate thymic chimeras, thymic lobes were obtained by dissection from neonatal Balb/c, Balb/B, and Balb/K mice and implanted into the interscapular dorsum of adult Balb/c *nu/nu* mice via a subcutaneous incision. Mice were maintained in sterile isolators on acidified water and rested 12 weeks prior to testing for chimerism and use in experiments.

10

### EXAMPLE 3

#### PREPARATION OF BLOOD-STAGE *P. FALCIPARUM* PARASITES

The FCB-1 line of *Plasmodium falciparum* were grown *in vitro* by the method of Trager and Jensen. Synchronous development of parasites was maintained by the sorbitol method of Lambros and Vanderberg. For the biosynthetic labelling of parasite proteins, <sup>3</sup>H-palmitic acid conjugated to defatted bovine serum albumin in molar ratio 1:1, <sup>3</sup>H-glucosamine or <sup>3</sup>H-mannose were added at a final specific activity of 10mCurie/ml, to RPMI 1640 cultures of 2x10<sup>10</sup> parasites at the late trophozoite/early schizont stage for 2 hours (for labelling of GPI precursors) or 8 hours (for labelling of protein-bound GPI). Parasites were harvested by 0.05% Saponin lysis and centrifugation in the cold at 15,000g for 20 minutes, followed by two washes in PBS and storage at -70°C.

25

### EXAMPLE 4

#### PURIFICATION OF THE 195KD MSP-1 AND 56KD MSP-2 ANTIGENS

The GPI-anchored MSP-1 and MSP-2 merozoite surface proteins were purified to homogeneity. Biosynthetically labelled malaria parasites at the late schizont stage were lysed in 0.05% Saponin and centrifuged at 15,000g for 20 minutes, and washed as above. The pellet was extracted in 25mM *n*-octyl-thioglucopyranoside (*n*-otg), 1%



- 65 -

BSA, 1mM EDTA, 0.1mM EGTA, 1mM PMSF, 1mM TPCK, 0.1mM TLCK, 5mM pCMPS, 1mg/ml pepstatin, 1mg/ml leupeptin, 1mM NEM, 5mM iodoacetamide, 150mM NaCl, 25mM Tris/HCl pH 7.4 by sonication on ice. The extract was clarified by centrifugation at 20,000g for 30 minutes in the cold, and the supernatant decanted and loaded onto two immunoaffinity columns arranged in sequence, containing approximately 10mg monoclonal antibody 111.4 or monoclonal antibody 113.1, each cross-linked to Protein G-Sepharose by glutaraldehyde (all procedures on ice). The protein extract was passed through the column at a rate of 0.3ml/min. The columns were washed first with 100ml 10mM n-otg, 1% BSA, 300mM NaCl, followed by 100ml 10mM n-otg, 300mM NaCl. Antigen was eluted from each column with four column volumes of 10mM n-otg, 200mM glycine pH 2.8. The pH of the eluate was neutralized with 2M Tris. Aliquots of protein were analysed for purity by SDS-PAGE followed by staining with Coomassie brilliant blue. The remaining purified proteins were dialysed exhaustively against 100mM  $\text{NH}_4\text{HCO}_3$  using dialysis membrane previously boiled exhaustively in 10mM EDTA followed by boiling in 10 changes of double distilled water. Protein concentration was determined by the method of Bradford.

#### EXAMPLE 5

#### 20 PURIFICATION OF MEMBRANE-FORM VARIANT SURFACE GLYCOPROTEIN (MF VSG) OF T. BRUCEI

*T. brucei* 118 (MITAT 1.5) was purified from the blood of infected Wistar rats by DEAE chromatography.  $1 \times 10^{11}$  parasites were pre-incubated for 30 minutes in glucose-deficient RPMI 1640 supplemented with 40mM fructose and then labelled in the same medium for 90 minutes either with [ $^3\text{H}$ ]-myristic acid conjugated to defatted BSA, or with [ $^3\text{H}$ ]-glucosamine. In the latter case the medium contained 1mg/ml tunicamycin. Parasites were washed in cold medium without label, and taken up in 10mM  $\text{ZnCl}_2$ , followed by centrifugation at 45,000g. The pellet was put into boiling 25mM n-otg, 5mM p-CMPS, 1mM PMSF, 1mM TLCK, 50mM Tris/HCL and allowed to cool, and centrifuged at 45,000g at 2°C. The remaining detergent soluble extract was made up to

- 66 -

1mM CaCl<sub>2</sub>, 1mM MgCl<sub>2</sub>, and 1mM MnCl<sub>2</sub>, and passed over a Con-A sepharose column, followed by washing with 10 column volumes of extraction buffer. The column was first eluted with detergent buffer containing 0.5M  $\alpha$ -methylmannopyranoside and 0.5M  $\alpha$ -glucopyranoside, followed by 25mM *n*-otg in 8M urea.

5 Aliquots were subject to SDS-PAGE and fluorography or staining with Coomassie blue.

### EXAMPLE 6

#### PURIFICATION OF THE C-TERMINAL GPI ANCHORS OF DEFINED PARASITE ANTIGENS

10 GPI-anchored *P. falciparum* MSP-1, MSP-2 and *T. brucei* 118 (MITat 1.5) mfVSG were labelled with fatty acid or glucosamine as required and purified as above, to 10mg/ml. 600ml methanol was added to 150ml aliquots followed by 150ml CHCl<sub>3</sub> and 450ml H<sub>2</sub>O. The samples were vortexed and microfuged, the supernatant taken for

15 scintillation counting, and the interphase and lower phase mixed with 450ml methanol and re-centrifuged. The pellet was repeatedly extracted with CMW 10:10:3 until partitioning of fatty-acid label into the supernatant was minimal, partitioned between water and water-saturated butanol, precipitated with acetone at -20°C, and the proteins taken up by sonication in 6M Urea, 1mM DTT, 1mM iodoacetic acid. After 15

20 minutes at room temperature, the sample was diluted 6 fold and made to 5mM CaCl<sub>2</sub>. 2.5% pre-digested Pronase B was added, and incubated for 72h at 37°C with 2 additions of 0.25% pronase. The sample was loaded in 5% 1-propanol, 0.1M NH<sub>4</sub>OAc onto pre-equilibrated Octyl-Sepharose, washed and eluted in a linear gradient of 1-propanol (5-60%) in water. GPIs eluted at 35-40% 1-propanol and were spotted onto

25 TLC plates (Si-60) and run in the solvent system C/M/HAc/W 25:15:4:2. The origin was scraped, GPIs eluted and partitioned between water and water-saturated butanol.

- 67 -

**EXAMPLE 7****PURIFICATION OF GIPLS AND GPI PRECURSORS BY TLC**

2x10<sup>3</sup>[H]-glucosamine-labelled *P. falciparum* schizonts were extracted in CM (2:1) and  
5 CMW (1:1:0.3), Folch washed, partitioned between water and water-saturated butanol,  
and dried. Residues were separated by 2D TLC (1st dimension CMW 4:4:1, 2nd  
dimension Butanol/HAc/W 4:6:1), the plates scanned by Bertold Digital  
Autoradiograph scanner, and the structurally defined GPI peaks scraped and re-  
extracted. Phospholipids were resolved away from GPI peaks. Areas lying outside the  
10 identifiable GPI peaks were treated in the same way, as were sham plates. GIPLs of *L.*  
*mexicana* were purified to homogeneity. Briefly, promastigotes were extracted twice in  
CMW (1:2:0.8), the insoluble material removed by centrifugation, and the CMW phase  
partitioned with 0.6 volume water. The dried upper phase was chromatographed on  
Octyl-Sepharose as above, and eluted GIPLs further purified by HPTLC using CM/1N  
15 NH<sub>4</sub>OH (10:10:3), and scrapings extracted with CMW (1:2:0.8). GIPL concentration  
and compositional purity was determined by GC-MS, following acid methanolysis and  
trimethylsilyl (TMS) derivatization. *myo*-Inositol content was measured following acid  
hydrolysis (6N HCl, 110°C, 16 h) and TMS derivatization, with selected ion  
monitoring for *m/z* 305 and 318. *scyllo*-Inositol was used as internal standard  
20 throughout.

**EXAMPLE 8****GENERATION OF CHEMICAL AND ENZYMATIC HYDROLYSIS  
FRAGMENTS OF GPIS AND GIPLS**

25 Purified, glucosamine-labelled *P. falciparum* and *T. brucei* GPIs, in which all dpms  
were detected in the organic phase following butanol/water partitioning, were subject to  
base hydrolysis by suspension in methanol/ammonia 1:1 for 6 hours at 50°C, followed  
by partitioning between water and water saturated butanol. Essentially 100% of label  
30 was then recovered from the aqueous phase. The aqueous phase was twice extracted  
with water-saturated butanol, lyophilized, and flash evaporated with methanol.

- 68 -

**EXAMPLE 9****DEAE ANION EXCHANGE CHROMATOGRAPHY**

GPIs were loaded onto a A DEAE column in 99% methanol, 1% water and washed  
5 with ten column volumes of solvent. They were subsequently eluted in 100mM  
Ammonium Acetate in 99% methanol, 1% water and dried under Nitrogen.

**EXAMPLE 10****BIOGEL P4 SIZE-EXCLUSION CHROMATOGRAPHY**

10

Base-hydrolysed GPI glycans were spiked with phenol red and blue dextran in 100mM  
Ammonium Acetate and further size-fractionated by passage through a 1cm x 1.2 metre  
Biogel P4 column equilibrated in 100mM Ammonium acetate in water. The column  
had previously been exhaustively calibrated by repeated analytical runs with GPI mixed  
15 with acid hydrolysed dextran markers to yield the relative elution position of glucose  
units detected by staining with orcinol in concentrated sulfuric acid. The column runs  
proved to be highly reproducible. For preparative purposes the dextran markers were  
omitted. The GPI peak was detected by scintillation counting of aliquots. Glycan  
concentration and compositional purity was determined by GC-MS, following acid  
20 methanolysis and trimethylsilyl (TMS) derivatization. *myo*-Inositol content was  
measured following acid hydrolysis (6N HCl, 110°C, 16 h) and TMS derivatization,  
with selected ion monitoring for *m/z* 305 and 318. *scyllo*-Inositol was used as internal  
standard throughout.

25

**EXAMPLE 11****COUPLING OF GPI GLYCAN TO MALEIMIDE-ACTIVATED PROTEINS**

GPIs were coupled to proteins by two methods. (i) GPIs were exposed to 1mM Traut's  
reagent (2-iminothiolane) in 40% 1-propanol, 60mM triethanolamine, 7mM potassium  
30 phosphate, 100mM NaCl, 1mM EDTA, pH 8.0 in the cold for 90 minutes under  
nitrogen. The sample was then desalted by gel filtration at 4°C through a small

- 69 -

immobilized dextran desalting column equilibrated in 40% 1-propanol in water. The propanol was dried off under nitrogen and the sample added to maleimide-activated KLH (Pierce) in coupling buffer (7mM potassium phosphate, 100mM NaCl, 1mM EDTA, pH 7.2) overnight followed by quenching with cysteine. (ii) (i) GPIs were  
5 exposed to 1mM Traut's reagent (2-iminothiolane) in 60mM triethanolamine, 30mM n-otg, 7mM potassium phosphate, 100mM NaCl, 1mM EDTA, pH 8.0 in the cold for 90 minutes under nitrogen. The sample was then desalted by gel filtration at 4°C through a small Biogel P4 column equilibrated in 30mM n-otg, 7mM potassium phosphate, 100mM NaCl, 1mM EDTA, pH 7.2 and added to maleimide-activated KLH (Pierce) in  
10 coupling buffer (7mM potassium phosphate, 100mM NaCl, 1mM EDTA, pH 7.2) overnight. The degree of conjugation was estimated by both by comparison of cpms before and after dialysis of the sample against PBS, or by use of Ellmans reagent to quantify sulfhydryl groups. Excess reactive groups were quenched with cysteine.

15

### EXAMPLE 12 FACS ANALYSIS

2x10<sup>5</sup> cells in ice cold murine tonicity RPMI 1640 with 0.05% Sodium azide and 1% BSA were incubated with optimally titrated FITC- , biotin- or phycoerythrin-conjugated  
20 monoclonal antibodies to murine CD3, CD4, CD5, CD8, CD25, CD28, CTLA-4, CD44, CD69, Thy-1 (NIMR-1 or G7), and NK1.1 as required. After washing in the same medium the cells were counter-stained with 0.5mg/ml propidium iodide and analysed by FACSScan. Cells were sorted on FACStar II.

25

### EXAMPLE 13 CYTOKINE CAPTURE ELISA ASSAYS

Cytokine levels were determined by specific capture ELISAs (Pharmingen). Treated and control samples were incubated at 37°C for 2 h in 96-well plates precoated with  
30 monoclonal anti-mouse IFN-g, IL-2 and IL-4, followed by washing and probing under the same experimental conditions with biotinylated second antibody anti-mouse IFN-g,

- 70 -

IL-2 and IL-4, followed by streptavidin-peroxidase. After addition of the substrate and color development, the plates were read by Titertek Multiscan MCC/340 automated ELISA reader. Cytokine mass was calculated by interpolating the results with the standard curve plotted with titrated recombinant mouse IFN- $\gamma$ , IL-2 and IL-4 of known mass.

#### EXAMPLE 14

#### ELISA ASSAY

10 Antigen (tetanus toxoid, *P. falciparum* rCS, *P. berghei* rCS, NANP<sub>40</sub>-BSA, FLU-BSA or BSA-alone) at 10ug/ml in phosphate binding buffer was incubated overnight in 50ul volumes in flat-bottomed Immulon 96-well plates, followed by extensive washing with buffer. The plates were blocked with 1% BSA in PBS for several hours. From a 1/32 dilution, sera were titrated two-fold in 1% BSA in PBS, and 50ul aliquots incubated in

15 triplicate for 2 hours at room temperature, followed by extensive washing with 1% BSA, 0.05% Tween-20 in PBS. An aliquot of affinity purified, biotin-labelled isotype specific goat anti-mouse second antibody was incubated as above, followed by further washing and the addition of streptavidin-alkaline phosphatase. After 30 minutes the plates were washed again and colourimetric development initiated by the addition of p-

20 Nitrophenylphosphate in diethanolamine buffer. Background binding to BSA-coated plates was determined in parallel. The titres derived are the last point giving values statistically different by two-way analysis of variance from non-specific binding by the same serum to the BSA-coated plates.

#### EXAMPLE 15

#### ANTIBODY FORMATION *IN VITRO*

Donor Balb/c *nu/nu* mice were primed twice with *P. berghei* SPZs or twice with LPS<sup>FLU</sup>. 10<sup>5</sup> splenocytes were placed in culture for 7 days in the presence of IL-2, with

30 and without antigen, anti-Class I, anti-Class II and anti-CD1 as indicated, with 10<sup>4</sup> CD4<sup>+</sup>, NK1.1<sup>+</sup> cells positively selected by FACS sorting and Dynal Detachabead from

- 71 -

spleens of C57Bl6 donors primed to GPI-KLH. *P. berghei* SPZs or GPI-OVA<sup>FLU</sup> were added with or without antibodies to CD1, MHC Class I or MHC Class II at various concentrations and cultured in RPMI 1640 + 10% FCS,  $5 \times 10^{-5}$  M b-ME and left for 8 days. Antigen-specific IgG production in the culture supernatant was determined by  
 5 ELISA against recCS and fluoresceinated Dog serum albumin (Sigma) as capture antigen.

### EXAMPLE 16

#### MHC II RESTRICTION IN THE ANTIBODY RESPONSE

10 This study determined that the antibody response to the native CS protein of malaria sporozoites is not Class II-restricted. Allogeneic bone-marrow irradiation chimeras, demonstrably unable to respond to the protein antigens Tetanus toxoid or recCS, were essentially equal to syngeneic chimeric controls in responding to irradiated sporozoites  
 15 with the production of anti-CS IgG1 and IgG2 (Table 1). Nude mice cannot respond to sporozoites with anti-CS IgG, and as reported previously (3), passive transfer of cytolytic anti-CD4, but not anti-CD8, antibodies into euthymic animals abrogated anti-CS antibody formation in response to sporozoites. However, nude mice engrafted with irradiated neonatal allogeneic thymi produced anti-CS IgG1 and IgG2 in response to  
 20 sporozoites at levels similar to recipients of syngeneic thymic implants, but did not respond to recCS or other protein antigens (Table 1). Class II<sup>0/0</sup> mice, lacking both Class II and Class II-restricted CD4<sup>+</sup> T cells, and reportedly incapable of mounting IgG responses to T-dependent protein antigens (4-6), responded to native but not recombinant CS protein with high titres of IgG1, IgG2a and IgG2b, similar to wildtype  
 25 controls (Table 2). Together, these data demonstrate that thymically-derived CD4<sup>+</sup> cells are required for IgG formation in response to the native CS protein, but that the Class II MHC is not a restriction element in this response.

- 72 -

### EXAMPLE 17

#### REQUIREMENT FOR GPI LIPID DOMAIN

To determine whether the GPI anchor could account for the difference in immunological behaviour of the native and recombinant proteins, GPIs purified from *Plasmodium falciparum*, *Leishmania mexicana* and *Trypanosoma brucei* were covalently linked in a molar ratio of 1:1 with haptenated Ovalbumin (GPI-OVA<sup>FLU</sup>). In contrast to sham-OVA<sup>FLU</sup> alone, GPI-OVA<sup>FLU</sup> was able to induce anti-FLU and anti-OVA IgG1, IgG2a and IgG2b formation in MHC Class II<sup>o/o</sup> mice (Table 2). IgG responses in Class II<sup>o/o</sup> mice were obtained only in response to OVA<sup>FLU</sup> coupled to intact GPI, and not to OVA<sup>FLU</sup> coupled to the deacylated GPI glycan, indicating the GPI lipid domain is required, and the glycan not sufficient, for the phenomenon (Table 2). However, GPI-OVA<sup>FLU</sup> was unable to induce anti-FLU IgG responses in nude mice or euthymic animals treated with lytic anti-CD4 indicating that the GPI does not provide a sufficient signal to drive immunoglobulin class switch in the absence of CD4<sup>+</sup> cells. Isotype distributions similar to those obtained by immunization of Class II<sup>o/o</sup> mice with GPI-OVA<sup>FLU</sup> were observed following exposure to the intact membrane form Variant Surface Glycoprotein of *T. brucei*, but not the deacylated soluble VSG derived by phospholipase C hydrolysis, confirming the requirement for GPI lipid domain and demonstrating that the phenomenon is generalizable to other GPI anchored proteins.

### EXAMPLE 18

#### PROLIFERATIVE AND FUNCTIONAL RESPONSES TO GPI ANCHORED PROTEINS

The *in vitro* proliferative and functional responses to purified GPIs of splenocytes from wild-type and Class II<sup>o/o</sup> animals primed with either neo-GPI-anchored proteins, *Plasmodium falciparum* or *P. berghei* sporozoites, and blood stage *P. chabaudi* infection were examined. In all cases, compositionally pure GPIs (by GC/MS) elicited proliferative responses from splenocytes. The blastoid cells responding to GPI were predominantly NK1.1<sup>+</sup>, CD4<sup>int</sup>, TCR $\alpha/\beta$ <sup>int</sup>, V $\alpha$ 14<sup>+</sup> and CD69<sup>+</sup>, and the FACS



- 73 -

profiles revealed both a relative and absolute increase in this cell population. This was also accompanied by the production of high levels of IL-4. This unusual population of T cells was considerably expanded following exposure of mice to sporozoites. The proliferative and cytokine response of NK1 + CD4+ T cells to purified GPIs could be blocked by anti-CD1 monoclonal antibody 1B1(7).

### EXAMPLE 19 ANTIBODY PRODUCTION

10 When splenocytes from sporozoite primed animals were exposed to sporozoites, they produced measurable levels of anti-CS IgG in the culture supernatant after 8 days. This antibody formation *in vitro* could be substantially blocked by anti-CD1 mAb 1B1, but less so by antibodies to Class II and not at all by anti-Class I. Furthermore, the NK1.1<sup>+</sup> cells were able to cooperate with B cells in CD1-restricted antibody formation  
15 to GPI-OVA<sup>FLU</sup> but not OVA<sup>FLU</sup>.

Wild-type and CD1<sup>o/o</sup> mice were exposed to sporozoites. Approximately 8-16 fold difference in anti-CS end-titres were detected, indicating that antibody formation to the native protein was reduced by approximately 90% in the latter animals, but in response  
20 to either TT or recCS was identical to wild-type controls (Table 3).

### EXAMPLE 20

COOH-terminal GPI's were purified from affinity-purified GPI-anchored proteins of *P. falciparum* (PfGPI) and *Trypanosoma brucei* membrane-form variant surface  
25 glycoprotein (mfVSG), and non-protein-linked free GPIs from *Leishmania mexicana*. The compositional purity of these latter molecules was confirmed by gas chromatography-mass spectrometry (GC-MS). In addition, a phosphorylated and lipidated mammalian GPI based on the rat brain Thy-1 GPI, and the corresponding  
30 inositolphosphoglycan (IPG) lacking a lipid tail, both chemically synthesized by n-pentenyl glucoside strategy and compositionally pure by <sup>1</sup>H NMR analysis, were also

- 74 -

used.

NK1.1<sup>+</sup> CD4<sup>+</sup> cells from wild-type and class II<sup>-/-</sup> mice, when cultured with irradiated wild-type or class II<sup>-/-</sup> antigen-presenting cells (APCs), responded to these purified GPIs as determined by incorporation of [<sup>3</sup>H]-thymidine ([<sup>3</sup>H]TdR), and the production of high levels of IL-4 and IFN- $\gamma$ . No proliferation in the absence of APCs indicated that GPIs do not provide a direct activation signal to NKT cells sufficient to induce cell growth. NKT cells did not respond to GPIs when cultured with irradiated APCs from  $\beta_2$ M<sup>-/-</sup> and CD1.1/CD1.2<sup>-/-</sup> (CD1<sup>-/-</sup>) donors, or with wild-type and class II<sup>-/-</sup> APCs in the presence of anti-CD1.1, but responded fully in the presence of isotype controls. The proliferative and IL-4 response to GPI of NKT cells and the V $\alpha$ 14<sup>+</sup>, CD4<sup>+</sup> subset in unfractionated splenocytes could also be blocked by the anti-CD1 mAb 1B1. Thus the recognition of GPIs by NKT cells is MHC-independent and CD1-restricted.

In addition, NKT cells produced IL-4 in response to CD1.1-transfected J774 macrophages in the absence of exogenous antigen, but not to sham-transfected controls. Nonetheless, the response was enhanced when CD1.1-transfectants were pulsed with GPI. However, in the absence of exogenous antigen no cytokine expression was detected in response to APCs expressing normal level of CD1.1. Thus high levels of CD1.1 expression in transfected cells may alone be sufficient to drive proliferation. Alternatively, as NKT cells can respond to mammalian GPIs (for example, synthetic Thy-1 GPI), CD1.1-transfectants may be able to present endogenous GPIs to NKT cells.

25

#### EXAMPLE 21

Both murine and human cerebral malaria are TH1-driven pro-inflammatory diseases. The *P. berghei* ANKA murine cerebral malaria model has several features in common with the human cerebral malaria syndrome. Specifically, it is a TNF- $\alpha$  and IFN- $\gamma$  dependent encephalitis associated with upregulation of ICAM-1 on the cerebral microvascular endothelium, an increase in both parasite and macrophage/neutrophil

- 75 -

adherence to these target cells, and attendant neurological complications. Unlike human cerebral malaria, there is a breakdown of the blood-brain barrier in the terminal stages of the murine syndrome. However, in the proximal stages the murine disease reflects more accurately the inflammatory cascade leading to cerebral involvement in humans. This syndrome manifests most clearly in C57B16 and CBA mice, and is clearly a IFN- $\gamma$  dependent, TH1 type disease. The TH2-promoting cytokine IL-4 can counter this condition and lack of IL-4 makes the condition worse.

Although susceptible to infection with *P. berghei*, Balb/c mice are usually refractory to the cerebral malaria syndrome (they preferentially manifest a TH2 response). It was observed however that Balb/c CD1.1/CD1.2 gene targeted (knockout) mice showed a high rate of *P. berghei*-induced cerebral malaria (Figure 1). This suggests that a CD1-dependent mechanism protects against cerebral malaria. Furthermore, *in vivo* lysis of NKT cells by passive transfer of lytic monoclonal antibody causes expression of the cerebral syndrome, demonstrating a protective effect against cerebral malaria of NKT cells. Consistent with these observations, it was shown, using intracellular cytokine staining and FACS analysis, that the cytokine profiles of conventional T cells in CD1-knockout mice were heavily biased to the TH1 pole i.e. there was a higher frequency of IFN- $\gamma$  producing cells (Figure 2). This indicates that CD1-dependent T cells play an important role in regulation of downstream TH1/TH2 differentiation. Thus priming against GPI in a CD1-dependent manner could inhibit the development of cerebral malaria in mice and humans.

## EXAMPLE 22

### RESULTS

T cell-dependent IgG responses to protein antigens are thought to be exclusively MHC class II- restricted. However, allogeneic bone-marrow irradiation chimeras were similar to syngeneic controls in responding to malaria sporozoites (SPZ) with IgG to the Circumsporozoite (CS) protein, despite inability to respond to the nominal protein antigens tetanus toxoid (TT) or a full-length recombinant *Plasmodium falciparum* CS

- 76 -

protein (recCS). Nude mice cannot respond to SPZ with anti-CS IgG, and passive transfer of depleting anti-CD4 antibodies into euthymic animals abolishes the anti-CS response to SPZ. However, nude mice engrafted with irradiated neonatal allogeneic thymi mounted anti-CS IgG responses to SPZ similar to recipients of syngeneic thymi, but did not respond to recCS or TT. Mice lacking both class II and class II-restricted CD4<sup>+</sup> T cells, and unable to respond to T-dependent antigens, produced anti-CS IgG (mean log<sub>2</sub> reciprocal titer of 10) in response to SPZ. Thus CD4<sup>+</sup> T cells are required for the IgG response to the native CS protein, but this may proceed through a MHC class II-independent route.

To determine whether the GPI anchor accounts for the difference in immunological behaviour of the proteins, we purified COOH-terminal GPIs from affinity-purified GPI-anchored proteins of *P. falciparum* (PfGPI) and *Trypanosoma brucei* membrane-form variant surface glycoprotein (mfVSG) (Fig. 3A), and non-protein-linked free GPIs from *Leishmania mexicana* (Fig. 3B). The compositional purity of these latter molecules was confirmed by gas chromatography-mass spectrometry (GC-MS). In addition, a phosphorylated and lipidated mammalian GPI based on the rat brain Thy-1 GPI (Fig. 3C), and the corresponding inositolphosphoglycan (IPG) lacking a lipid tail, both chemically synthesized by *n*-pentenyl glucoside strategy and compositionally pure by <sup>1</sup>H NMR analysis, were also used. To generate responses to the hapten fluorescein (FLU), native and synthetic GPIs and IPG were exposed to 2-iminothiolane to introduce a sulfhydryl onto free amino groups, de-salted and conjugated in a molar ratio of 1:1 to fluoresceinated, maleimide-activated ovalbumin (OVA<sup>FLU</sup>). In contrast to sham-OVA<sup>FLU</sup> alone, or sham-OVA<sup>FLU</sup> mixed with equal molar amounts of free PfGPI, PfGPI-OVA<sup>FLU</sup> conjugates were able to induce anti-FLU IgG1 formation in MHC class II<sup>-/-</sup> mice (mean log<sub>2</sub> reciprocal titer of 8). Similar IgG responses were obtained in class II<sup>-/-</sup> mice with the mfVSG of *T. brucei*, but not the deacylated sVSG derived by PI-PLC hydrolysis (Fig. 3A), demonstrating that the GPI lipid domain is required, and the GPI glycan is not sufficient, for the phenomenon. This was confirmed by comparing responses to OVA<sup>FLU</sup> conjugated to either synthetic Thy-1 GPI or Thy-1 IPG lacking fatty acid (log<sub>2</sub> reciprocal titer of 9.75 vs. no response respectively). Thus IgG

- 77 -

responses in class II<sup>-/-</sup> mice require linkage of antigen to GPI with an intact lipid, which may be of diacylglycerol or alkylacylglycerol composition.

The *in vitro* proliferative and cytokine responses to purified malarial GPI of splenocytes from animals primed with malaria SPZ was examined. There was a marked increase in both the relative and absolute numbers of NK1.1<sup>+</sup> CD4<sup>+</sup> blastoid cells responding to GPI from both class II<sup>-/-</sup> (Fig. 4A) and wild-type mice. A high frequency of both V $\alpha$ 14<sup>+</sup> CD4<sup>+</sup> (Fig. 4A) and V $\beta$ 8<sup>+</sup> cells was also detected in the responding population. No exogenous cytokines were required for this proliferation, but supplementation of cultures with 5U/ml IL-2 increased the level of response.

To examine the fine specificity of responding cells, splenocytes from wild-type and class II<sup>-/-</sup> animals primed to *P. berghei* SPZ were exposed to 0.5 $\mu$ M of the structures shown in Fig. 3, plus dipalmitoyl-PI. The cells responded to a similar degree to most intact GPIs, but only weakly to the iM2 GPI with truncated glycan and not at all to glycans lacking the fatty acid domain, or to PI (Fig. 4B). Thus both glycan and fatty acids are required for recognition, and NKT cells from SPZ-primed donors respond to a range of GPIs from diverse protozoal and mammalian taxa. However, as the full range of structures presented to the host under these priming conditions is not known, the results may reflect either broad recognition of diverse antigens by the general population of NKT cells or heterogeneous responses from a clonally mixed population. To analyze this further, cells were expanded in the presence of either PfGPI or Thy-1 GPI for four days, rested in IL-2, then re-stimulated with homologous or heterologous antigen. Cells expanded by either antigen responded significantly less well to the heterologous stimulus (Fig. 4B). Analysis of antigen-specific frequencies following *in vitro* culture revealed no increase above background in response to PfGPI in naïve donors, but a clear increase in antigen-reactive NKT cells from SPZ-primed donors (7-30% of total). However, naïve and primed donors both mounted significant responses to the iM4 GPI (to 5% of NKT cells).

30

- 78 -

When sorted NK1.1<sup>+</sup> CD4<sup>+</sup> cells from wild-type and class II<sup>-/-</sup> mice primed to SPZs were cultured with irradiated wild-type or class II<sup>-/-</sup> antigen-presenting cells (APCs), they responded to purified GPIs as determined by incorporation of [<sup>3</sup>H]-thymidine ([<sup>3</sup>H]TdR), and the production of high levels of IL-4 (Fig. 5A). No proliferation in the absence of APCs indicated that GPIs do not provide a direct activation signal to NKT cells sufficient to induce cell growth. NKT cells did not respond to GPIs when cultured with irradiated APCs from  $\beta_2M^{-/-}$  and CD1.1/CD1.2<sup>-/-</sup> (CD1<sup>-/-</sup>) donors, or with wild-type and class II<sup>-/-</sup> APCs in the presence of anti-CD1.1 (1B1), but responded fully in the presence of isotype controls (Fig. 5A). The proliferative and IL-4 response to PfGPI of NKT cells and the V $\alpha$ 14<sup>+</sup>, CD4<sup>+</sup> subset in unfractionated splenocytes could also be blocked by the anti-CD1 mAB 1B1 (Fig. 5B). Thus the recognition of GPIs by NKT cells in MHC-independent and CD1-restricted. In addition, NKT cells produced IL-4 in response to CD1.1-transfected J774 macrophages in the absence of exogenous antigen, but not to sham-transfected controls. Nonetheless, the response was enhanced when CD1.1-transfectants were pulsed with PfGPI (Fig. 5A). The response of NKT cells to CD1.1, observed previously, has been adduced in support of the proposition that this cell population may play a physiological role in the absence of associative recognition of antigen. However, as reported, in the absence of exogenous antigen no cytokine expression was detected in response to APCs expressing normal level of CD1.1. Thus high levels of CD1.1 expression in transfected cells may alone be sufficient to drive proliferation. Alternatively, as NKT cells can respond to mammalian GPIs (for example, synthetic Thy-1 GPI (Fig. 4B)), CD1.1-transfectants may be able to present endogenous GPIs to NKT cells.

Extending to antigen-specific systems, NKT cells cooperated with B cells by ELISPOT assay in CD1-restricted IgG formation to GPI-OVA<sup>FLU</sup> and native *P. berghe* CS protein, but not OVA<sup>FLU</sup> (Fig. 6A). To determine therefore whether CD1.1 or CD1.2-restricted antibody formation was a major or minor contributor to the IgG response to GPI-anchored proteins and SPZ *in vivo*, CD1<sup>-/-</sup> mice and wild-type controls were exposed to mfVSG<sup>FLU</sup>, SPZ or recCS. Responses to mfVSG<sup>FLU</sup> and SPZ were significantly curtailed in CD1<sup>-/-</sup> mice (Fig. 6B), indicating that under these conditions

- 79 -

the CD1-restricted pathway of IgG formation is a significant component of responses to the native CS protein. Both groups responded equally to recCS, confirming that class II-dependent responses are unaffected by loss of CD1.

## METHODS

Irradiated H-2 congenic mice received  $10^7$  T-depleted syngeneic or allogeneic bone-marrow cells. Thymic lobes from neonatal Balb/c, Balb/B, and Balb/K mice were irradiated and implanted into adult Balb/c *nu/nu* mice. Mice were rested 12 weeks prior to testing for chimerism. *P. falciparum* and *P. berghei* SPZ were dissected from *Anopheles freeborni* and *A. stephensi* respectively. Some preparations were kept at  $-70^\circ\text{C}$  before use. Mice were primed with TT, recCS, or SPZ, boosted after 1 week, and IgG responses analyzed by ELISA. RecCS, derived from the T4 strain and consisting of the entire sequence except for the C-terminal 21 amino acids (GPI-anchor signal sequence), was used as immunizing antigen and to detect responses to *P. falciparum* SPZs. The *P. berghei* rCS (detection antigen only for *P. berghei* SPZs) encompasses amino acids 81-277. Antigen-specific end-titers were defined as the last titration giving values statistically different from binding to plates without antigen. A  $\log_2$  reciprocal titer of 4.0 was the background cutoff. To TT, recCS and *P. berghei* SPZ, syngeneic thymic chimeras mounted mean  $\log_2$  reciprocal IgG titers of 15.14, 13.5 and 13.5, respectively. The equivalent values for allogeneic chimeras were 4.5, 4.25 and 13. For syngeneic bone-marrow chimeras the equivalent values were 15.11, 11.25 and 14.6, and for allogeneic chimeras 8.5, 7.25 and 15.25, respectively. The results were confirmed using *P. falciparum* SPZ.

IgG responses to rCS, recCS or FLU were determined as above in class II<sup>-/-</sup> and wild-type mice receiving *P. berghei* SPZ, recCS, mfVSG<sup>FLU</sup>, sVSG<sup>FLU</sup> or OVA<sup>FLU</sup>, either sham-conjugated (sham-OVA<sup>FLU</sup>) or conjugated to *P. falciparum* GPI (PfGPI-OVA<sup>FLU</sup>), Thy-1 GPI (Thy-1-OVA<sup>FLU</sup>) or Thy-1 IPG (IPG-OVA<sup>FLU</sup>). Unlike wild-types, Class II<sup>-/-</sup> animals failed to raise IgG to recCS, sVSG<sup>FLU</sup>, sham-OVA<sup>FLU</sup> and IPG-OVA<sup>FLU</sup> (mean  $\log_2$  reciprocal titers < 4). However, they produced IgG in response to SPZ, PfGPI-

- 80 -

OVA<sup>FLU</sup>, mfVSG<sup>FLU</sup> and Thy-1-OVA<sup>FLU</sup> (log<sub>2</sub> reciprocal titer of 10,8 8.25 and 9.75, respectively). Ig isotypes included IgG1, IgG2a and IgG2b.

- mfVSG and GPI-anchored *P. falciparum* proteins, purified by HPLC and affinity chromatography were extracted in chloroform:methanol, precipitated with acetone, solubilized and reduced, diluted in 5mM CaCl<sub>2</sub>, digested for 72h at 37°C with Pronase B, purified over Octyl-Sepharose, partitioned between water and water-saturated butanol (PfGPI), and purified by high performance thin layer chromatography (HPTLC) ( $R_f=0.05$ ) using C/M/HAc/W (25:15:4:2). CMW (1:2:0.8) extracts of *L. mexicana* promastigotes were purified over Octyl-Sepharose followed by HPTLC using CM/1N NH<sub>4</sub>OH (10:10:3). The compositional purity of the latter molecules was confirmed by GC-MS following acid methanolysis and trimethylsilyl (TMS) derivatization for neutral monosaccharide analysis, and acid hydrolysis (6N HCl, 110°C, 16 h) and TMS derivatization, with selected ion monitoring for  $m/z$  305 and 318. *scyllo*-Inositol was used as internal standard throughout. Compositional molar ratios were as predicted and no contaminating carbohydrates or sugars were detected. The specific values are published elsewhere as are the <sup>1</sup>H NMR compositional analyses demonstrating purity of the synthetic Thy-1 GPI and IPG.
- Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations of any two or more said steps or features.



- 81 -

Table 1

MHC Class II is not a restriction element in IgG formation to SPZs.

Test group	Antigen:	TT	recCS	nativeCS
BM chimeras				
Balb/c----- > Balb/c				
	Animal#101	32768	32768	ND
	Animal#103	65536	4096	ND
	Animal#114	32768	8192	ND
	Animal#107	131072	32768	ND
	Animal#106	16384	ND	65536
	Animal#111	32768	ND	32768
	Animal#105	32768	ND	16384
	Animal#121	8192	ND	32768
	Animal#119	65536	ND	8192
Balb/c----- > Balb/K				
	Animal#202	512	64	ND
	Animal#203	128	128	ND
	Animal#205	1024	256	ND
	Animal#207	128	256	ND
	Animal#208	256	ND	32768
	Animal#211	512	ND	16384
	Animal#212	256	ND	65532
	Animal#217	1024	ND	65532
Thymic chimeras				
Balb/c----- > Balb/c				
	Animal#1006	65536	6384	ND
	Animal#1007	16384	16384	ND
	Animal#1014	131072	4096	ND
	Animal#1016	16384	ND	16384
	Animal#1022	32768	ND	8192
	Animal#1025	131072	ND	16384
	Animal#1028	8192	ND	8192
Balb/c----- > Balb/K				
	Animal#2004	32	64	ND
	Animal#2005	16	32	ND
	Animal#2006	< 16	< 16	ND
	Animal#2010	128	< 16	ND
	Animal#2014	32	ND	4096
	Animal#2019	32	ND	16384
	Animal#2020	16	ND	8192
	Animal#2022	< 16	ND	8192

- 82 -

Table 2.

IgG responses to the recCS, native CS, OVA<sup>FLU</sup> and GPI-OVA<sup>FLU</sup>  
in C57 wild-type, Class II<sup>-/-</sup> and  $\beta_2$ M<sup>-/-</sup> mice.

Primed with:	Reciprocal titre to:			
	CS		FLU	
	SPZs	recCS	OVA <sup>FLU</sup>	GPI-OVA <sup>FLU</sup>
1) Wild-type	4096	4096	4096	8192
	4096	8192	1024	512
	4096	8192	2048	512
	16384	16384		< 16
2) Class II <sup>-/-</sup>	512	< 16	< 16	128
	512	< 16	< 16	64
	2048	< 16	< 16	512
	2048	32	32	1024
3) $\beta_2$ M <sup>-/-</sup>	1024	8192	ND	ND
	2048	8192		
	2048	8192		
	4096	8192		

- 83 -

Table 3.

The IgG response to the native CS protein is reduced in CD1.1<sup>0/0</sup> mice

Immunized with:	Reciprocal anti-CS titre	
	SPZs	recCS
1) Balb/c	4096	4096
	4096	8192
	8192	8192
	16384	16384
2) Balb/c CD1 <sup>0/0</sup>	512	8192
	512	8192
	1024	8192
	2048	8192

- 84 -

**BIBLIOGRAPHY:**

1. McConville, M.J. and Ferguson, M.A.J. *Biochem. J.* 294:305 (1993).
2. Stevens, V.L. *Biochem. J.* 310:361 (1995).
3. Weiss, W.R., Sedegah, M., Berzofsky, J.A. and Hoffman, S.L. *J. Immunol.* 151:2690 (1993).
4. Cosgrove, D. *et al.*, *Cell* 66:1051 (1991).
5. Grusby, M.J., Johnson, R.S., Papaioannou, V.E. and Glimcher, L.H. *Science* 253:1417 (1991).
6. Köntgen, F., Süss, G., Stewart, C., Steinmetz, M. and Bluethmann, H. *Int. Immunol.* 5:957 (1993).
7. Bleicher, P.A. *et al.*, *Science* 250:679 (1990).